School of Public Health

Thesis and Dissertations

2022-01

# Central Obesity and Associated factors Among Adults in Debre Tabor Town, Northeast, Ethiopia, 2021: A Community-Based Cross-Sectional Study

Tigist, Yismaw

http://ir.bdu.edu.et/handle/123456789/13629 Downloaded from DSpace Repository, DSpace Institution's institutional repository



# BAHIR DAR UNIVERSITY COLLEGE OF MEDICINE AND HEALTH SCIENCES DEPARTMENT OF NUTRITION AND DIETETICS

# CENTRAL OBESITY AND ASSOCIATED FACTORS AMONG ADULTS IN DEBRE TABOR TOWN, NORTHEAST, ETHIOPIA, 2021: A COMMUNITY-BASED CROSS-SECTIONAL STUDY

BY TIGIST YISMAW (BSc, PUBLIC HEALTH)

A THESIS SUBMITTED TO THE DEPARTMENT OF NUTRITION AND DIETETICS, COLLEGE OF MEDICINE AND HEALTH SCIENCES SCHOOL OF PUBLIC HEALTH, BAHIR DAR UNIVERSITY IN PARTIAL FULFILMENT OF THE REQUIREMENTS FOR THE DEGREE OF MASTER OF PUBLIC HEALTH NUTRITION

> JANUARY 2022 BAHIR DAR, ETHIOPIA

# BAHIR DAR UNIVERSITY COLLEGE OF MEDICINE AND HEALTH SCIENCES SCHOOL OF PUBLIC HEALTH DEPARTMENT OF NUTRITION AND DIETETICS

CENTRAL OBESITY AND ASSOCIATED FACTORS AMONG ADULTS IN DEBRE TABOR TOWN, NORTHEAST, ETHIOPIA, 2022: A COMMUNITY-BASED CROSS-SECTIONAL STUDY.

# PRINCIPAL INVESTIGATOR: TIGIST YISMAW (BSc IN PUBLIC HEALTH) Tel: +251918280918

Email: tgyismaw21@gmail.com

ADVISORS:

Dr NETSANET F. (MPH, DOUBLE PHD, ASSOCIATE PROFESSOR OF HUMAN NUTRITION)

Tel: +251913516677

E-mail: netsanetfentahun01@gmail.com

Ms HANNA D. (MPH IN NUTRITION, ASST. PROFESSOR)

Tel: +251913360728

Email: hanna.de21@gmail.com

A THESIS SUBMITTED TO THE DEPARTMENT OF NUTRITION AND DIETETICS, SCHOOL OF PUBLIC HEALTH, COLLEGE OF MEDICINE AND HEALTH SCIENCES IN THE PARTIAL FULFILMENT OF THE REQUIREMENTS FOR THE DEGREE OF MASTERS IN PUBLIC HEALTH NUTRITION

> JANUARY, 2022 BAHIR DAR, ETHIOPIA

> > I

# Acknowledgement

First of all, I would like to thank the almighty GOD for giving me the strength and diligence to complete this study.

Secondly, my heartfelt special thanks extend to my respected advisors Dr Nestanet F. and Ms Hanna D. for their invaluable advice and comment to do this research thesis.

Furthermore, I extend my thanks to Debre Tabor town administrators and for all the data collectors. As well, my special thanks extend to my friends and colleagues.

Last but not least, I thank you Bahir Dar University and Simada Woreda Health office for the chance to give me this MPH degree opportunity at Bahir Dar University.

#### Abstract

**Background**: Central obesity nowadays is becoming a chronic non-communicable disease that is an indicator of mortality and morbidity. In Ethiopia obesity becomes a major public health problem. As far as we are concerned, there is no evidence in the study area on the prevalence of central obesity and factors affecting central obesity.

**Objective**: To assess the magnitude of central obesity and its associated factors among adults aged 18-64 years in Debre Tabor Town, northeast, Ethiopia, 2021.

**Methods**: A community-based cross-sectional study was conducted. A total of 616 study participants were included in this study. Systematic random sampling was employed to select the study participants. Waist circumference was measured to declare central obesity. The level of physical activity was measured using the global physical activity questionnaire (GPAQ). Depression, Anxiety and Stress Scale (DASS-21) was used to screen depression among adults. Data were entered using EpiData and analyzed using STATA/MP version 16 software. Bivariable and multivariable logistic regression was employed to identify factors affecting central obesity. A P-value less than 0.05 was used to declare statistical significance.

**Results**: A total 582 study participants recruited with a response rate of 94.48%. The overall prevalence of central obesity among adults in Debre Tabor town 37.80% (95% CI: 33.94 – 41.82). Higher prevalence of obesity was observed among females (55.08 %) than male adults (26.01%). Female sex (Adjusted odds ratio (AOR) = 5.81, 95% Confidence interval (CI): 3.01 - 11.20), aged 18 - 35 years (AOR = 3.84, 95% CI: 2.30 - 6.41), aged 56-64 years (AOR = 5.90, 95% CI: 2.76 - 12.64), mild depressed (AOR = 9.72, 95% CI: 3.99 - 23.64), and physical inactive (AOR = 1.58, 95% CI:1.01 - 2.51) were had high odds for central obesity; while daily labor workers (AOR =0.43, 95% CI: 0.20 - 0.91) was low odds for central obesity.

**Conclusion:** More than one-third of adults in Debre Tabor town had central obesity. Physically inactive and mild depression were modifiable factors for central obesity. Age and sex were non-modifiable factors for central obesity. Individuals and policymakers should give prior attention to modifiable factors for central obesity. Health promoters and policymakers should give primary attention for females.

Keywords: Central Obesity, Central Obesity, Physical Activity, Adults, Ethiopia.

# Table of contents

Cor	Contents Pages			
Ack	nowle	edger	ment	II
Abs	tract.			111
Tab	le of o	conte	ents	IV
Abb	revia	tions	and acronyms	VI
List	of tal	oles		VII
List	of fig	ures.		.VIII
1.	Intro	oduct	tion	I
1	.1.	Back	kground	I
1	.2.	State	ement of the problem	2
1	.3.	Sign	ificance of the study	4
2.	Liter	ature	e Review	5
2	.1.	The	magnitude of central obesity	5
2	.2.	Fact	ors associated with central obesity	6
	2.2.2	1.	Socio-demographic factors	6
	2.2.2	2.	Behavioral factors	6
	2.2.3	3.	Dietary factors	7
3.	Con	ceptu	ual Framework	8
4.	Obje	ective	25	9
4	.1.	Gen	eral objective	9
4	.2.	Spec	cific objectives	9
5.	Met	hods	and materials	10
5	.1.	Stud	dy setting and period	10
5	.2.	Stud	dy design	10
5	.3.	Рор	ulations	10
	5.3.2	1.	Source populations	10
	5.3.2	2.	Study populations	10
5	.4.	Eligi	bility criteria	10
	5.4.3	1.	Inclusion criteria	10
	5.4.2	2.	Exclusion criteria	10
5	.5.	Sam	ple size determination and sampling procedure	10

	5.	5.1.	Sample size determination		
	5.	5.2.	Sampling technique and procedure		
	5.6.	Vari	ables of the study		
	5.	6.1.	Outcome variable		
	5.	6.2.	Independent Variables		
	5.7.	Ope	rational definitions	14	
	5.8.	Data	a collection tools and procedures	14	
	5.9.	Data	a quality control	16	
	5.10	. D	ata management and analysis	16	
	5.11	. Et	thical consideration	16	
6.	Re	esults			
	6.1.	Soci	o-demographic and behavioral characteristics of the respondents		
	6.2.	Foo	d consumption status among study participants		
	6.3.	Prev	valence of central obesity		
	6.4.	Fact	ors associated with central obesity	20	
7.	Di	iscussio	n	23	
8.	Co	onclusio	on	25	
9.	<ol> <li>Recommendations</li></ol>				
1(	).	Refere	nces	27	
11	L.	Annex			
12	2.	Amhar	ic version questionnaire		
13. Declaration		Declar	ation	43	

# Abbreviations and acronyms

DASS	Depression Anxiety Stress Scale
FFS	Food Consumption Score
GOPA	Global Observatory for Physical Activity
GPAQ	Global Physical Activity Questionnaire
LMICs	Low- and Middle-Income Countries
METs	Metabolic Equivalents Tasks
NCDs	Non-Communicable Disease
STATA	Statistics and Data
WC	Waist Circumference
WHO	World Health Organization
WHR	Waist Hip Circumference
WHtR	Waist to Height Ratio

# List of tables

Table 1: Socio-demographic and behavioural characteristics of the study participants, 2021	17
Table 2: Food consumption status in a typical week among adults in Debre Tabor town, 2021	19
Table 3: Magnitudes of central obesity among adults in Debre Tabor Town, 2021	20
Table 4: Multivariable logistic regression analysis factors associated with central obesity among adult	ts in
Debre Tabor town, 2021.	21

# List of figures

Figure 1: Conceptual framework for central obesity adapted from deferent literature [17,45,48,51,52]	8
Figure 2: Sampling procedure to select study participants	13

#### 1. Introduction

#### 1.1. Background

Central obesity is a medical condition defined as abnormal or excessive fat accumulation (regionally, globally, or both) which increases the health risk [1]. It is a health risk that increases body weights and fat distributions that lead to the expression of co-morbid diseases that occur at different thresholds depending on the population [2,3]. Obesity classification has different measurements. It would be based on a practical measurement widely available and easily practical depending on their setting. But the most accurate prediction of health risk (prognosis) measures of body fat (the major component of body weight responsible for adverse outcomes) such as underwater weighing, dual-energy x-ray absorptiometry (DEXA) scanning, computed tomography (CT), and magnetic resonance imaging (MRI) are impractical for use in everyday clinical encounters. Estimates of body fat including body mass index (BMI) and waist circumference do have limitations compared to these imaging methods, but still, provide relevant information and are easily implemented in a variety of practice settings [2]. In practice, according to the World Health Organization (WHO) central or truncal obesity is defined by waist circumference (WC) [4], which is WC  $\geq$  94 cm for men and  $\geq$  80 cm for women were a cut point to declare central obesity. The International Diabetes Federation (IDF) puts different cut-off points for different ethnic groups such as 94 cm for males and 80 cm for females for Europeans, 90 cm for males and 80 cm for females for Asians) [4].

Based on excess body fat distribution obesity is classified into three. The first one is the general fat type of obesity in which the whole body from head to toe looks like a tub. The second is the Pear type in this type, the lower part of the body has the extra flesh. This type of obesity is more common for females. The flesh is somewhat flabby in the abdomen, thighs, buttocks and legs. The third, Android/Central type of obesity is likened to the shape of an apple. The shoulders, face, arms, neck, chest and upper portion of the abdomen are bloated. The stomach gives a stiff appearance. The lower portion of the body is thinner beyond proportion in comparison with the upper part. Though this type of obesity is found more in males, it is also present to a lesser extent in women. Android type of obesity is a major risk for heart damage and heart disease due to high cholesterol [5].

#### **1.2.** Statement of the problem

Central obesity nowadays is becoming a chronic non-communicable disease that is an indicator of mortality and morbidity [2]. It is also a problem in low- and middle-income countries [6]. World health organization (WHO) puts obesity as the fifth leading cause of mortality globally and it is one of the chief health challenges [7]. Worldwide, central obesity has nearly tripled since 1975 and in 2016, more than 1.9 billion and 650 million adults, older than aged 18 years, were overweight and obese, respectively [3]. Even though central obesity and overweight is a problem of high-income countries; low- and middle-income countries (LMICs), in particular urban settings of sub-Saharan Africa (SSA), face the challenge of an increasing trend [8]. Out of the 20 fast-rising countries with adult central obesity, nearly half of them are found in Africa [9].

The burden of obesity is higher among women and urban residents. On the other hand, previous studies evidenced that the risk of central obesity is multifactorial. Socioeconomic status, age, parity, marital status, physical inactivity, and increased energy are powerful predictors of central obesity in SSA [10]. Physical inactivity during the transition from adolescence to adulthood is related to overall obesity and severe central obesity in females [11]. Physical inactivity is the most important risk factor for the rising prevalence of obesity [12]. The number of people who are obese is increasing due to modern lifestyle changes, physical inactivity, and excessive consumption of energy-dense foods [13,14]. The odds of maintaining a healthy body weight significantly increased with engagement in > 500 MET per week (approximately 167 min per week) [15]. Adhering to vigorous-intensity physical activity of at least 1 h per week was associated with a significant decrease in the incidence of obesity [16].

Few studies in Ethiopia showed that a significant number of individuals were centrally obese. A study in Northwest, Ethiopia, Dilla town Ethiopia, Gondar and Woldiya town Ethiopia showed that 16.5% to 37.6% [17–20]. World Health Organization (WHO) recommends the reduction of obesity through healthy eating and physical activity [21]. The sustainable development goal (SDG) 3, target 3.4 calls to reduce by one-third premature mortality related to non-communicable diseases through prevention and treatment and promote mental health and well-being [22].

Globally obesity is a major public health problem including low and middle-income countries. The prevalence of overweight and obesity has increased significantly in the nutritional transition period, and the health burden of obesity-related complications is growing. The introduction of fast and westernized dietary habits is the mainstay of the reason for the increasing prevalence of obesity. A few years ago, countries were in a problem with under-nutrition but due to nutrition transition and sedentary behavior obesity becomes a major public health challenge and a double burden of Undernutrition especially in SSA. The cause of the non-communicable disease (NCDs) is mostly modifiable and can be preventable if diagnosed earlier. As one of the main modifiable risk factors for NCDs was obesity.

Different nutritional programs in Ethiopia [23,24] gives prior attention to undernutrition which undermines the alarming rise of obesity due to unhealthy diet. There is a paucity of evidence in the study area about the magnitude of central obesity and associated factors. Therefore, this study aimed to assess the level of central obesity and associated factors among adults aged 18-64 years at Debre Tabor Town, northeast, Ethiopia.

#### **1.3.** Significance of the study

Measuring central obesity is a key role in the diagnosis of this metabolic-related disease. Mostly the common method to measure obesity at community level body mass index. The finding of the study was helpful to encourage local people to improve current nutritional status by handling the risk factors and improving feeding pattern of healthy diet and physical activity. Encourage government and other stakeholders for the development of programs and policies related to nutrition based on the facts and figures generated by this study. Documentation of such information will provide insight into policy intervention to tackle the alarming increase of central obesity in Ethiopia.

Hence, the result or outcome of this study was providing the valuable information for researcher, policy makers, NGOs; governmental and non-governmental organizations as well as local administrations to plan future circumstances.

### 2. Literature Review

#### 2.1. The magnitude of central obesity

A study conducted on a Spanish adult population revealed that the prevalence of central obesity was 33.4% of the higher prevalence observed among women (43.3%) than men (23.3%) [25]. A cross-sectional survey conducted at Madurai, Tamil Nadu, India showed that the magnitude of central obesity was 20.9% among adults [26]. A study conducted in rural Bangladeshi showed that the age-standardized prevalence of central obesity was 39.8% of this the prevalence of central obesity was higher than among women than men [27].

Different cross sectional studies conducted in Africa evidenced that the prevalence of central obesity ranges from 11.8% to 67.8% [28–35]. The prevalence of central obesity in Africa was higher than among women (19.5% - 39.2%) than men (1.3% - 13.8%) [32–35].

A study conducted in Woldia town, Northeast Ethiopia revealed that the prevalence of central obesity was 16.5% (95% CI: 14.2– 19.2) which is higher than women (27.9%) than men (3.9%) [20]. In a study conducted by University employees in Ethiopia, the magnitude of waist circumference is 33.6% [19]. An institutional-based cross-sectional study in Ethiopian university employees showed that 33.6% of adults were abdominally obese [36]. Another study in urban areas of Northwest Ethiopia showed that central obesity among adults was 37.6% [17]. Furthermore, a community-based cross-sectional study in Dilla town, Ethiopia revealed that 24.4% of adults were abdominally obese and a higher prevalence was observed among women adults [18]. Generally, there are limited study on the prevalence of central obesity, even if some studies are available, there is a tool, sample size, and sociodemographic difference among inference population. In addition to this some of the variables significantly associated were inconsistent among literatures. Therefore, this study aimed to assess the level of central obesity and associated factors among adults aged 18-64 years at Debre Tabor Town, northeast, Ethiopia.

#### 2.2. Factors associated with central obesity

#### 2.2.1. Socio-demographic factors

A meta-analysis report among 13.2 million subjects showed that older adult individuals, female individuals, and urban residents had a statistically significant association with central obesity [37]. A study from Spain indicates that, males and aged older than 40 years were associated with a higher risk of central obesity, whereas those who had a higher level of education were associated with a lower risk of central obesity [38].

A study conducted from India, Tanzania, Eastern Sudan, and Uganda [31,33,39,40] revealed that females were more likely develop central obesity than males. Studies from India and Uganda showed that urban residents are also more at risk of developing central obesity than rural residents [33,39]. Previous studies from Uganda, Tanzania, and Eastern Sudan revealed that married or cohabiting had a higher odds of central obesity than their counterparts [31,33,40].

In a study conducted in India showed that higher socio-economic status, were a significant determinant factor for central obesity [39]. A study from Tanzania showed that age greater than 30 years were more likely to develop central obesity than younger adults [40].

A study in urban areas of Northwest Ethiopia showed that with a one-year increase in age the odds of developing central obesity would increase by 5%. As well this study revealed that female sex was a risk factor for central obesity [17]. On the other a community-based cross-sectional in Woldia town, Northeast Ethiopia showed that female sex, higher household wealth class, age from 35 to 55 years, aged older than 55 years, secondary educational level were risk factors for central obesity, whereas single marital status decreased the central obesity [20]. A study in Dilla town showed that abdominally obese were observed among adults with a high and middle house wealth class, and being female adults [18].

#### 2.2.2. Behavioral factors

Different studies revealed that behavioral factors are associated with central obesity. A study in Spain showed that current smokers were 28% less likely to develop central obesity. On the other hand, individuals who had spent more than 75 minutes per week physical activity were less likely to develop central obesity [38]. Another study in China showed that alcohol drinking and former cigarette smokers were more likely centrally obese than their counterparts [41]. A longitudinal study evidenced that physical activity significantly lowers central obesity [42]. A population-based study from 31 countries revealed that ex-smokers were positively associated

with obesity, while light or moderate smoker were inversely related with central obesity among both men and women adult populations. This study also showed that moderate alcohol use was positively associated with obesity, while heavy alcohol use was negatively associated [43].

A study in Switzerland showed that current smokers had lower mean waist circumference compared with non-smokers [44]. In a study from Spain showed that individuals who had spent more than seven hours per day in sleeping was associated with a lower risk of central obesity [38]. A study in Brazil showed that physical activity either aerobic or resistance exercises, with a prolonged duration of moderate-intensity for men and walking or moderate intensity for women is suggested as the most adequate for the prevention strategies of central obesity [45]. A systematic review and meta-analysis review showed that exercise had a greater effect on the outward appearance of obesity (body mass index, waist circumference) than on its practical factors [46]. A study from Dilla town, Ethiopia showed that being physically inactive were more at risk to develop central obesity [18]. A study conducted in Jimma, Ethiopia showed that khat chewers had a lower body mass index and fat-free mass as compared to non-chewers [47].

#### 2.2.3. Dietary factors

A longitudinal study showed that dietary fiber intake (increased fruit and low carbohydrate) significantly decreased the occurrence of central obesity [42]. A study from a sample of 31 countries indicates that the daily consumption of fruit and vegetables was found to be protective against central obesity [43]. Another study showed that poor dietary patterns had direct associations with central obesity only in men [48]. Previous studies evidenced that eating out of home was increased the risk of central obesity [49]. A study showed that high consumption of energy-rich diet increases excess body fat accumulation. This study also evidenced that bad eating habits such as calorie-rich foods, skipping breakfasts, drinking sugar-sweetened beverages, eating out frequently, and emotional eating increases the risk of central obesity [50].

# **3.** Conceptual Framework

The Conceptual frameworks of central obesity are listed under the diagram.



Figure 1: Conceptual framework for central obesity adapted from deferent literature [17,45,48,51,52].

# 4. Objectives

#### 4.1. General objective

To assess the prevalence of central obesity and its associated factors among adults aged 18-64 years in Debre Tabor Town, Northeast, Ethiopia, 2021.

# 4.2. Specific objectives

To determine central obesity among adults aged 18-64 years in Debre Tabor Town, Northeast, Ethiopia, 2021.

To identify associated factors for central obesity among adults aged 18-64 years in Debre Tabor Town, Northeast, Ethiopia, 2021.

# 5. Methods and materials

#### 5.1. Study setting and period

Debre Tabor town is the central town of South Gondar Zone, Amhara National Regional State. The town is located in northeast Ethiopia, 667 km away from Addis Ababa the capital city of Ethiopia. The city is found at approximately 11.64° latitude and 37.90° with an elevation of 2,706 meters above sea level. According to the 2007 national census report, the town has a total population of 55,596 populations, of whom 27,644 are men and 27,952 women. Debre Tabor has 6 kebeles with a total of 15, 549 households. The study was conducted from August/22/2021 to September/10/ 2021.

# 5.2. Study design

A community-based cross-sectional study was conducted.

# 5.3. Populations

# 5.3.1. Source populations

Among Adults aged 18 -64 years living in Debre Tabor town.

# 5.3.2. Study populations

Among Adults living in Debre Tabor town in the selected kebeles (01 & 06) during the data collection period.

# 5.4. Eligibility criteria

# 5.4.1. Inclusion criteria

Among Adults aged 18-64 years and living more than at least six months before the data collection period were included in this study.

# 5.4.2. Exclusion criteria

Pregnant women were excluded from this study.

# 5.5. Sample size determination and sampling procedure

# 5.5.1. Sample size determination

Sample size determination was calculated based on the objectives of the study as follows: Based on the first objective using Epidemiologic information version 7 software to calculate sample size by taking prevalence of the outcome using the single population proportion formula.

The formula is

$$n = \frac{(Z_{\alpha/2})^2 * p(1-p)}{d^2}$$

Where n = the minimum sample size required for this study

 $Z_{a/2} = Z$ - value at the alpha value of 0.05 which is 1.96

d = margin of error which is taken as 4%

P = Prevalence of central obesity from the previous study.

To calculate the minimum sample size using the single population proportion formula a study conducted at Dabat Health and Demographic Surveillance Site (HDSSs) in Gondar [17] was appropriate instead of almost similar demography and recent study. In this study, the prevalence of central obesity is 37.6. But the prevalence is less than 50% the margin of error should be less than 5% which was an appropriate 4% to make representativeness. Assuming the above formula the final sample size was 616 after adding a 10% non-response rate.

Based on the second objective the sample size was calculated as follows using Kelsey [53] formula.

 $n1 = \frac{(Z_{\frac{\alpha}{2}} + Z_{1-\beta})^2 p' q'(r+1)}{r(p1-p2)^2} \quad , \qquad n2 = r * n1, \quad p' = \frac{p1+r*p2}{r+1} \quad , \text{ and } q' = 1-p'$ 

Where

n1 = number of exposed (non-reference category)

n2 = number of unexposed (reference category)

 $Z_{\alpha/2} = z$ -score for a two-tailed test based on  $\alpha$  level (5% level)

 $Z_{1-\beta}$  = z-score for a one-tailed test based on  $\beta$  level (80% power)

r = ratio of reference group vs non reference group

p1 = proportion of central obesity among exposed

p2 = proportion of central obesity among unexposed

Based on the above formula the different sample size was calculated using different factors by Epi-Info version 7 software. Therefore, based on the above study in different factors.

Factors

Assumption

Required sample size

Sex (male reference group) [17]	$r = 0.62 \ \Box \ 1, p1 = \ 52\%, \ p2 = 13.8\%$	46
Type of dietary oil (solid reference) [17]	$r = 2.29 \Box 2, p1 = 57.9\%, p2 = 28.8\%$	99
Physical activity (active reference) [18]	$r = 2.5 \Box 3, p1 = 39.2\%, p2 = 18.5\%$	178

Therefore, the sample calculated using a single population proportion is the best representative.

#### 5.5.2. Sampling technique and procedure

To made representativeness 30% of among the six Kebele, two Kebeles were selected using a lottery method; while kebele 01 and kebele 06 were selected. All are urban kebeles. The two kebeles had a total number of 5261 households. Systematic random sampling techniques was employed in those kebeles in the 9<sup>th</sup> interval. House to house survey was employed in systematically selected households. Marking of the household was done to avoid double selection.

Finally, one eligible adult was selected from the selected household and if more than one found is eligible adultin .the household the Kish grid sample size selection method was applied [54].



Figure 2: Sampling procedure to select study participants.

#### 5.6. Variables of the study

#### 5.6.1. Outcome variable

Central / abdominal obesity (Y/N)

# 5.6.2. Independent Variables

Scio-Economic factors:	Educational status
Age	Occupation
Sex	Household wealth index
Religion	<b>Behavioral factors:</b>
Marital status	

Smoking statusPhysical activity statusAlcohol consumptionPhysical activity statusKhat chewingDepression, anxiety, and stress statusDiet-related factors:Food consumption scoreFood consumption scorePoor food consumptionBorderline food consumptionAcceptable food consumptionAcceptable food consumptionEating out of home

#### 5.7. Operational definitions

**Central obesity:** Waist circumference (WC)  $\ge$  94 cm for males and  $\ge$  80 cm for females according to WHO definition [4].

**Physically active:** Individuals achieve moderate and vigorous-intensity physical activity achieving at least 600 metabolic equivalent tasks (MET) minutes [55].

**Physically inactive:** Individuals achieve moderate and vigorous-intensity physical activity achieving less than 600 metabolic equivalent tasks (MET) minutes [55].

**Food consumption status:** Individuals had a sum score was calculated from the eight food items using 0 - 21, 21.5 - 35, and > 35 poor, borderline, and good food consumption status, respectively [56].

**Depression, anxiety, and stress status:** Normal (0-9), mild (10-13), moderate (14-20), severe (21-27), and extremely severe (28+) depression was declared according to their respective scores. Normal (0-7), mild (8-9), moderate (10-14), severe (15-19), and extremely severe (20+) anxiety scale was declared and normal (0-14), mild (15-18), moderate (9-25), sever (26-33), extremely severe (34+) stress scale was declared [57].

#### **5.8.** Data collection tools and procedures

The data were collected using a structured interviewer-administered questionnaire. The data were collected by four clinical nurses by house-to-house survey. Central obesity or central obesity was measured between the anterior superior iliac span and the lower coastal margin perpendicular to the midclavicular line using a non-stretchable tap meter. During measurement, the closes were removed to get the true measurement.

#### **Socioeconomic factors**

We measured socio-demographic and economic-related factors found to be associated with the central obesity structured questionnaire. Household wealth index status was measured in five categories (poorest, poor, middle, rich, and richest) after the household questionnaire survey questionnaire.

#### **Behavioral related factors**

A structured interviewer-administered questionnaire was prepared from WHO-STEP wise for noncommunicable diseases. Physical activity was assessed using the global physical activity questionnaire (GPAQ) developed by the WHO for physical activity surveillance [55]. The Global Physical Activity Questionnaire was developed by WHO for physical activity surveillance in countries. It collects information on physical activity participation in three domains as well as sedentary behavior, comprising 16 questions (P1-P16). The domains are categorized as vigorous work-related, vigorous recreational, moderate work-related, moderate recreational and recreational physical activity. The questionnaire assesses physical activity in the past seven days paraphrasing as in a typical week, on how many days do participate in physical exercise. The level of physical activity was in hours and minutes on a typical day in a week. Finally, the questionnaire was finalized in metabolic equivalent per task (METs) to classify physically active and inactive.

Depression, anxiety, and stress were assessed using the Depression, Anxiety and Stress Scale - 21 scale (DASS-21) is a set of three self-report scales designed to measure the emotional states of depression, anxiety and stress. Each of the three DASS-21 scales contains 7 items, divided into subscales with similar content. The depression scale assesses dysphoria, hopelessness, devaluation of life, self-deprecation, lack of interest/involvement, anhedonia and inertia. The anxiety scale assesses autonomic arousal, skeletal muscle effects, situational anxiety, and subjective experience of anxious affect. The stress scale is sensitive to levels of chronic nonspecific arousal. It assesses difficulty relaxing, nervous arousal, and being easily upset/agitated, irritable / over-reactive and impatient. Scores for depression, anxiety and stress are calculated by summing the scores by multiplying two for the relevant items [57].

#### **Dietary related factors**

Dietary consumption of oil type, eating out of home and food consumption status was measured. The food consumption score (FCS) dietary consumption assessment tool was prepared by the United nation World food program [56]. The food consumption score (FCS) is a composite score based on dietary diversity, food frequency, and relative nutritional importance of different food groups. Food items are grouped into 8 standard food groups (cereals and tubers, pulses, vegetables, fruit, meat and fish, milk, sugar and Oil) with a maximum value of 7 days per week. Sum score was done to calculate individualize score of the study participants. In the FCS the highest score is a good dietary habit.

#### 5.9. Data quality control

The data quality was assured during tool development, data collection, coding, entry and analysis. A pretest was done before the date of data collection at kebele 04. The training was delivered for data collectors before the data collection period. The questionnaire was prepared in English then translated to Amharic again back to translate into English for its consistency. Waist circumference was measured in a non-stretchable tap meter and a double measurement was made. From two measurements the average value was taken.

#### 5.10. Data management and analysis

The data were checked, coded, and enter in Epi-data version 4.6 software and it was imported in STATA/MP version 16 statistical software for further analysis. Descriptive statistics were reported in tables and graphs. The principal component analysis was employed to determine the household wealth index. The chi-square assumption was a check for the model assumption. Bivariable logistic regression was carried out. Two sides p-value less than 0.25 was used to select candidate variables for multivariable logistic regression. Hosmer and Lemeshow's goodness of fit test was checked for model fitness. Multicollinearity was checked using the variance inflation factor (VIF < 10). Two sides p-value less than 0.05 was considered to declare statistical significance. Finally adjusted odds ratio with a 95% confidence level was reported.

#### 5.11. Ethical consideration

Ethical clearance was secured from the institutional review board (IRB) of Bahir Dar University. Letter permission was also secured from Debre Tabor City Administration. Informed consent was secured from each study participant. Hand warm was made during waist circumference measurement. Participants' identifiers were omitted from the questionnaire and respondents can withdraw at any time if they do not want to participate and data was kept confidential and anonymous.

# 6. Results

#### 6.1. Socio-demographic and behavioral characteristics of the respondents

Overall, a total of 582 study participants were recruited with a response rate of 94.48%. The mean  $\pm$  SD age of the study participants was 36.10  $\pm$  11.90 years. The majority (334, 57.39%) of the study participants were below the age group of 35 years. Fifty nine percent of the study participants were males. Almost all, (571, 98.11%) of the study participants were Christian Orthodox in their religion. Half of the respondents had college and above level of education. A total of 196 (33.68%) of study participants were in poor household wealth status. Overall, 39% of the respondents drunk standard alcohol (Vodka, Areki...) preceding one year of the survey. Of the total of respondents, 15 (2.59%) and 24 (4.12%) had a history of the current smoker and chat chewer, respectively. Moreover, 45.02% and 20.10% of the study participants were moderately and vigorously physically active, respectively. Overall, a total of 300 (51.55%) of study participants were physically active (**Table 1**).

Variables	Categories	Frequency (n)	Percentage (%)
Sex	Male	346	59.45
	Female	236	40.55
	Young adults (18–35 years)	334	57.39
	Middle-aged adults (36–55 years)	196	33.68
Age	Older adults (55 - 64 years)	52	8.93
Marital status	Union	391	67.18
	Not-union	191	32.82
Religion	Orthodox Christian	571	98.11
	Others §	11	1.89
	No education	57	9.79
	Primary	103	17.70
Educational status	Secondary	134	23.02
	College and above	288	49.48
Occupational status	Government employ	220	37.80
	Merchant	91	15.64
	Daily Labor	98	16.84

Table 1: Socio-demographic and behavioral characteristics of the study participants, 2021. (n = 582)

	Housewife	118	20.27
	Student	55	9.45
Household wealth	Poor	196	33.68
status	Middle	147	25.26
	Rich	239	41.07
Alcohol use	No	113	19.42
	Yes	469	80.58
Standard alcohol use	No	355	61.00
	Yes	227	39.00
Current chat chew	No	558	95.88
	Yes	24	4.12
Current smoker	No	565	97.41
	Yes	15	2.59
Ever smoker	No	526	90.38
	Yes	56	9.62
Moderate physical	Inactive	320	54.98
activity	Active	262	45.02
Vigorous physical	Inactive	465	79.90
activity	Active	117	20.10
Physical activity status	Inactive	282	48.45
	Active	300	51.55
Total		582	100

NB: § = Muslim and Catholic

#### 6.2. Food consumption status among study participants

Overall, 261 (44.85%) had borderline food consumption status and 277 (47.59%) acceptable food consumption status. The mean  $\pm$  SD of food consumption score was 36.96  $\pm$  12.44; the minimum and the maximum 7.5 and 82, respectively. Among the food items vegetables, pulses, cereals, and oils are the most consumed food in the typical week, while fruit and milk are the least consumed food (**Table 2**).

-	Response		
Food item	Yes (%)	Mean per	
		day	
In a typical week, on how many days do you eat fruit?	272 (46.74)	0.58	
In a typical week, on how many days do you eat vegetables?	522 (89.69)	2.20	
In a typical week, on how many days do you eat protein source foods	321 (55.15)	1.08	
from animals (beef, lamb, chicken, fish, and egg)?			
In a typical week, on how many days do you eat protein source foods	575 (98.80)	4.91	
from plants (pea, bean, chickpea, nuts, groundnuts)?			
In a typical week, on how many days do you eat energy source foods	548 (94.16)	3.45	
(cereal grains, energy-rich tubers such as potatoes, sweet potatoes)?			
In a typical week, on how many days do you eat milk and milk	156 (26.80)	0.75	
products (milk, cheese, Yogurt)?			
In a typical week, on how many days do you eat fats (fats and oils)?	577 (99.14)	6.74	
In a typical week, on how many days do you eat /drink discretionary	368 (63.23)	2.37	
calories (soft drinks, sugar, chocolates, honey)?			
Over all Food Consumption Score (FCS)			
$\checkmark$ Poor food consumption = 44 (7.56%)			
$\Im$ Borderline food consumption = 261 (44.85%)			

Table 2: Food consumption status in a typical week among adults in Debre Tabor town, 2021.

NB: Food items in the yes category had at least one time per week otherwise no.

#### 6.3. Prevalence of central obesity

race Acceptable food consumption = 277 (47.59%)

The overall prevalence of central obesity among adults in Debre Tabor town was 37.80% (95% CI: 33.94 - 41.82). Females 130 (55.08%) were more obese than males 90 (26.01%) and central obesity was higher in older-aged adults. Central obesity was higher among physically inactive individuals 133 (47.16%). Moreover, central obesity was higher among depressed adults (**Table 3**).

		Central Obesity s	tatus
Variables		Not obese n (%)	Obese n (%)
Sex of the respondent	Male	256 (73.99)	90 (26.01)
	Female	106 (44.92)	130 (55.08)
Age	Young adults	256 (75.15)	83 (24.85)
	Middle-aged adults	90 (45.92)	106 (54.08)
	Older adults	21 (40.38)	31 (59.62)
FCS	Poor food consumption	31 (70.45)	13 (29.55)
	Borderline food consumption	171 (65.52)	90 (34.48)
	Good food consumption	160 (57.76)	117 (42.24)
Depression status	Normal	333 (72.23)	128 (27.77)
	Mild	10 (12.20)	72 (87.80)
	Sever	19(48.72)	20 (51.28)
Anxiety status	Normal	342 (68.13)	160 (31.87)
	Mild	5 (17.86)	23 (82.14)
	Sever	15 (28.85)	37 (71.15)
Stress status	Normal	309 (65.47)	163 (34.53)
	Mild	41 (65.08)	22 (34.92)
	Sever	12 (25.53)	35 (74.47)
Physical activity status	Inactive active	149 (52.84)	133 (47.16)
	Active	213 (71 )	87 (29)
Central obesity		362 (62.20)	220 (37.80)

Table 3: Magnitudes of central obesity among adults in Debre Tabor Town, 2021

#### 6.4. Factors associated with central obesity

In bi-variable logistic regression analysis sex, age, marital status, occupation status, eating out of home, ever smoking status, household wealth status, type of oil use, depression, stress, and physical activity were statistically significant factors for central obesity. In multivariable logistic regression analysis sex, age, occupation status, depression, and level of physical activity were statistically significant with central obesity.

Keeping all other variables constant, the odds of central obesity among females were almost six times higher than males (AOR = 5.81, 95% CI: 3.01 - 11.20). The odds of developing central

obesity among individuals aged 36-55 years were four times more likely than younger adults aged (18 to 35 years) (AOR = 3.84, 95% CI: 2.30 - 6.41). Individuals aged 56-64 years were six times more likely to develop central obesity as compared to younger-aged adults (AOR = 5.90, 95% CI: 2.76 - 12.64). Daily labor workers were 57% less likely to develop central obesity as compared to government employees (AOR = 0.43, 95% CI: 0.20 - 0.91). The odds of developing central obesity among mild depressed adults were 9.72 times higher than their counterparts (AOR = 9.72, 95% CI: 3.99 - 23.64). Furthermore, physically inactive individuals were 58% more likely vulnerable for central obesity as compared to physically active individuals (AOR = 1.58, 95% CI: 1.01 - 2.51) (**Table 4**).

Table 4: Multivariable logistic regression analysis factors associated with central obesity among adults in Debre Tabor town, 2021.

	Central obesity				
Independent variables	Yes	No	COR (95%CI)	AOR (95%CI)	
Sex					
Female	130	106	3.49 (2.45 – 4.96)	5.81 (3.01 – 11.20) ***	
Male	90	256	1	1	
Age in years					
Old adults	21	51	4.46 (2.43 - 8.19)	5.90 (2.76 – 12.64) ***	
Middle age adults	106	90	3.56 (2.45 - 5.18)	3.84 (2.30 - 6.41) ***	
Young adults	83	251	1		
Marital status					
Not Union	46	145	0.40 (0.27 – 0.58)		
Union	174	217	1		
Occupational status					
Daily labor	20	78	0.40 (0.23 – 0.70)	0.43 (0.20 – 0.91) *	
Merchant	40	51	1.22 (0.75 – 2.00)	1.33 (0.71 – 2.48)	
House wife	65	53	1.91 (1.22 – 3.00)	0.49 (0.24 – 1.00)	
Student	9	46	0.30 (0.14 – 0.65)	0.61 (0.22 – 1.67)	
Government employ	86	134	1		

Food consumption status				
Good	117	160	174 (0.87 – 3.48)	
Borderline	90	171	1.26 (0.63 – 2.52)	
Poor	13	31	1	
Meals eat out of home				
Yes	49	150	0.40 (0.28 - 0.59)	
No	171	212	1	
Ever smoker				
Yes	105	46	0.33 (0.16 – 0.66)	
No	210	316	1	
Household wealth status				
Rich	79	160	0.62 (0.42 - 0.91)	
Middle	106	90	0.73 (0.47 – 1.13)	
Young adults	83	251	1	
Oil use				
Solid oil	147	73	0.68 (0.47 - 0.98)	
Liquid oil	271	91	1	
Depression status				
Sever	20	19	2.74 (1.42 - 5.29)	1.38 (0.55 – 3.48)
Mild	72	10	18.73 (9.37 – 37.42)	9.72 (3.99 – 23.64) ***
Normal	128	333	1	
Stress status				
Sever stress	37	8	10 (4.59 – 22.16)	
Mild stress	23	5	10 (5.75 – 26.87)	
Normal	160	349	1	
Physical activity				
Inactive	133	149	2.19 (1.55 - 3.08)	1.58 (1.01 – 2.51) *
Active	87	213	1	

NB: \*\*\* = Significant at 0.001 level, \*\* = Significant at 0.01 level, \* = Significant at 0.05 level, COR = Crude Odds Ratio, AOR = Adjusted Odds Ratio, CI = Confidence Interval.

#### 7. Discussion

Central obesity is becoming a major public health problem in the general population and it becomes increasing in low-middle income countries [58]. Central obesity is a common health condition related to a wide range of non-communicable diseases such as cardiovascular diseases, type-2 diabetes mellitus, and cancers. Nowadays it is an emerging nutritional transition health problem in low and middle-income countries, including Ethiopia. Therefore, documenting and conducting possibly modifiable risk factors will help policymakers to curve the burden of the problem.

This study revealed that the prevalence of central obesity among adults in Debre Tabor town was 37.80% (95% CI: 33.94 - 41.82). The prevalence of central obesity was higher among females (55.08 %) than males (26.01%). The finding was similar to previous studies conducted in Ethiopia 37.6% [17] and Iran (40.7%) [59]. This finding is lower than a study conducted in South Africa (67%) [28], West Africa (50.8%) [60], Nigeria (52.6%) [61], India (71.2%) [62], and China (52%) [30]; whereas, this study is higher than a study conducted in Ethiopia (16.5 – 24.4%) [18,20], Uganda (11.8%) [63], Iran (28.3%) [20] and Tanzania (24.88%) [29]. The possible source of lower or higher prevalence variation might be the deference central obesity definition cut points, setting deference, central obesity measurement deference and study period difference. Different studies might be use different cut point for waist circumference and some studies might conducted in only urban of rural setting. Therefore, the discrepancy might be due to those reasons.

This study showed that the likelihood of developing central obesity among females were six times higher than males. This finding agreed with a study conducted in different countries in the world [17,29,31,64–68]. This might be because of female sex had less physically active during their daily life. The physiological changes during reproductive ages may contribute to the change of their status of obesity [69–71]. Additional explanations for higher central obesity for females might be genetic, biological factors with females having more steroid hormones which expose them to obesity [66,72,73]. The other possible source of variation could be that in the Ethiopian cultural context males were more engaged in activities that require higher energy than females.

This study revealed that the age of individuals is significantly associated with central obesity. Individuals aged older than 35 years were more likely to develop central obesity as compared to adults younger than aged 35 years. This finding is supported by a study conducted in different countries [31,60,68,74,75]. This might be due to the reason that ageing is associated with insulin

resistance that changes body fat distribution and metabolism [74,75]. The other justification might be older age related to low physical activities and a tendency for a more sedentary way of life exposed to obesity. Daily labor workers were 57% less likely to develop central obesity as compared to government employees. The possible justification might be daily works might engage with vigorous physical activity.

This study also revealed that depression symptom was directly related to central obesity. Mild depressive adults were almost ten times more likely vulnerable to central obesity than their counterparts. A systematic review and meta-analysis study supports that the odds of central obesity among depressive individuals were higher by 38% than the reference category [45]. In addition, previous studies showed that central obesity and level of depression symptoms had a positive association [76–78]. Molecular and clinical studies revealed evidence for the association between depression and central obesity i.e. there is influence on hypothalamic pituitary adrenal (HPA) axis, inflammatory pathways and insulin sensitivity [79], which is abnormal regulation of HPA axis, the elevation of certain inflammatory markers and insulin resistance being individuals obese. Disturbance of the hormones can affect the secretion or metabolism of neurotransmitters, such as serotonin, norepinephrine and dopamine in the brain leading to central obesity which consequently influences mood disturbance.

Physical activity was statistically significant with central obesity. The odds of developing central obesity among physically inactive adults were 58% higher than physically active adults. Previous studies [17,18,80] had similar evidence with this study. Physical activity increases total energy expenditure, which can help them stay in energy balance and lose [81,82]. Therefore, physical activity decreases fat accumulation around the waist and total body fat, which decreases central obesity. Besides, physical activity reduces depression and anxiety and this might enhance physical exercise. Thus, aerobic and resistance exercises with a prolonged duration of moderate-intensity for men and walking or moderate intensity for women is suggested for the prevention or treatment of central obesity [45]. This study follows some limitations. Since the data were collected using an interview-based survey, it might have social desirability bias and recall bias for food consumption assessment. As well, the depression, anxiety, and stress screening might be affected since the data were collected during the instability period. Moreover, the limitation of the cross-sectional design is inevitable.

# 8. Conclusion

More than one-third of adults in Debre Tabor Town were abdominally obese. Females were more vulnerable than males for central obesity. Physical inactivity and depression among adults were modifiable risk factors for central obesity. Sex and age were non-modifiable risk factors for central obesity.

# 9. Recommendations

According to the evidence of this study, we recommend that the community itself and policymakers should give prior attention to preventable (primarily physical inactivity and depression) risk factors for central obesity.

### For the community:

They should practice regular physical activity to reduce central obesity as per the recommendations.

# For the policymakers and health planners

The policymakers and health planners should promote the benefit of physical activity to reduce central obesity.

Health promoters should give primary attention for those females which are at risk for central obesity.

Health promotion and primary prevention should promote the risks of central obesity on noncommunicable diseases.

# **10. References**

- 1. Purnell JQ. Definitions, classification, and epidemiology of obesity. Endotext [Internet]. 2018.
- 2. Purner JQ. Definitions, Classification, and Epidemiology of Obesity. Feingold KR, Anawalt B, Boyce A, al, Ed Endotext [Internet] South Dartmouth MDText.com, Inc; 2000-. 2018; 1–13.
- 3. WHO key fact. Obesity and Overweight. In: Encyclopedia of Adolescence [Internet]. 2021. doi:10.1007/978-1-4419-1695-2\_447
- 4. World Health Organisation (WHO). WHO | Waist Circumference and Waist–Hip Ratio. Report of a WHO Expert Consultation. Geneva, 8-11 December 2008. 2008. pp. 8–11. Available: http://www.who.int
- 5. S. AK, C. V. L. Prevalence and Risk Factors of Obesity among Women. Indian J Public Heal Res Dev. 2020;11: 437. doi:10.37506/v11/i2/2020/ijphrd/194840
- 6. World Health Orgnaisation key facts. Obesity. 2021. Available: https://www.who.int/news-room/facts-in-pictures/detail/6-facts-on-obesity
- 7. World Health Organisation (WHO). Report of the Commission on Ending Childhood Obesity. Implement Plan Exec Summ. 2017;(WHO/NMH/P: Licence: CC BY-NC-SA 3.0 IGO.
- 8. Biadgilign S, Mgutshini T, Haile D, Gebremichael B, Moges Y, Tilahun K. Epidemiology of obesity and overweight in sub-Saharan Africa: A protocol for a systematic review and meta-analysis. BMJ Open. 2017;7: 7–10. doi:10.1136/bmjopen-2017-017666
- 9. Mwakideu C. Obesity in Africa: A looming health catastrophe. Dw.Com. 2019. Available: https://www.dw.com/en/obesity-in-africa-a-looming-health-catastrophe/a-49920972
- 10. Charles Agyemang, Sandra Boatemaa, Grace Agyemang Frempong and A de-GA. Obesity in Sub-Saharan Africa. Metab Syndr. 2016; 447–459. doi:10.1007/978-3-319-11251-0
- Tammelin T, Laitinen J, Näyhä S. Change in the level of physical activity from adolescence into adulthood and obesity at the age of 31 years. Int J Obes. 2004;28: 775–782. doi:10.1038/sj.ijo.0802622
- 12. Weinsier RL, Hunter GR, Heini AF, Goran MI, Sell SM. The etiology of obesity: relative contribution of metabolic factors, diet, and physical activity. Am J Med. 1998;105: 145–150. doi:10.1016/S0002-9343(98)00190-9
- 13. Vandevijvere S, Chow CC, Hall KD, Umali E, Swinburn BA. Increased food energy supply as a major driver of the obesity epidemic: a global analysis. Bull World Health Organ. 2015;93: 446–456. doi:10.2471/blt.14.150565
- 14. Prentice AM. The emerging epidemic of obesity in developing countries. Int J Epidemiol. 2006;35: 93–99. doi:10.1093/ije/dyi272
- Brown WJ, Kabir E, Clark BK, Gomersall SR. Maintaining a Healthy BMI: Data From a 16-Year Study of Young Australian Women. Am J Prev Med. 2016;51: e165–e178. doi:10.1016/j.amepre.2016.09.007
- 16. Frankenberg E. Physical Activity and the Incidence of Obesity in Young African-American Women.

Bone. 2012;23: 1-7. doi:10.1016/j.amepre.2013.04.016.Physical

- Molla MD, Wolde HF, Atnafu A. Magnitude of central obesity and its associated factors among adults in urban areas of Northwest Ethiopia. Diabetes, Metab Syndr Obes Targets Ther. 2020;13: 4169–4178. doi:10.2147/DMSO.S279837
- Tesfaye TS, Zeleke TM, Alemu W, Argaw D, Bedane TK. Dietary diversity and physical activity as risk factors of abdominal obesity among adults in Dilla town, Ethiopia. PLoS One. 2020;15: 1–11. doi:10.1371/journal.pone.0236671
- 19. Janakiraman B, Abebe SM, Chala MB, Demissie SF. Epidemiology of general, central obesity and associated cardio-metabolic risks among university employees, Ethiopia: A cross-sectional study. Diabetes, Metab Syndr Obes Targets Ther. 2020;13: 343–353. doi:10.2147/DMSO.S235981
- 20. Dagne S, Menber Y, Petrucka P, Wassihun Y. Prevalence and associated factors of abdominal obesity among the adult population in Woldia town, Northeast Ethiopia, 2020: Community-based cross-sectional study. PLoS One. 2021;16: 1–16. doi:10.1371/journal.pone.0247960
- 21. Waxman A. WHO global strategy on diet, physical activity and health. Food Nutr Bull. 2004;25: 292–302. doi:10.1177/156482650402500310
- 22. WHO. World Health Organization | Noncommunicable diseases. Who. 2019. Available: https://www.euro.who.int/en/health-topics/health-policy/sustainable-developmentgoals/publications/2017/fact-sheets-on-the-sustainable-development-goals-sdgs-health-targets
- 23. Policy N. We shall strive to create food and nutrition self-reliant Ethiopia 01 Food and Nutrition Policy Federal Democratic Republic of Ethiopia. 2018.
- 24. FDR. Ethiopian National Nutrition Programme. Fed Democr Repub Ethiop. 2015. Available: https://extranet.who.int/
- Aranceta-Bartrina J, Pérez-Rodrigo C, Alberdi-Aresti G, Ramos-Carrera N, Lázaro-Masedo S. Prevalence of General Obesity and Abdominal Obesity in the Spanish Adult Population (Aged 25–64 Years) 2014–2015: The ENPE Study. Rev Española Cardiol (English Ed. 2016;69: 579–587. doi:10.1016/j.rec.2016.02.009
- 26. Easwaran M, Sivasubramanian P, Kannan G. Prevalence of central obesity and its association with sociodemographic profile among young adults attending Outdoor Patient Department of Community Health Centre in Madurai, Tamil Nadu. Int J Med Sci Public Heal. 2019;8: 1. doi:10.5455/ijmsph.2019.0719721092019
- Siddiquee T, Bhowmik B, Da Vale Moreira NC, Mujumder A, Mahtab H, Khan AKA, et al. Prevalence of obesity in a rural Asian Indian (Bangladeshi) population and its determinants. BMC Public Health. 2015;15: 1–9. doi:10.1186/s12889-015-2193-4
- 28. Owolabi EO, Ter Goon D, Adeniyi OV. Central obesity and normal-weight central obesity among adults attending healthcare facilities in Buffalo City Metropolitan Municipality, South Africa: a cross-sectional study. J Heal Popul Nutr. 2017;36: 54. doi:10.1186/s41043-017-0133-x
- Munyogwa MJ. The Prevalence of Abdominal Obesity and Its Correlates among the Adults in Dodoma Region, Tanzania: A Community-Based Cross-Sectional Study. Hindawi, Advances Med. 2018;2018: 8.
- 30. Mohamed SF, Haregu TN, Khayeka-Wandabwa C, Muthuri SK, Kyobutungi C. Magnitude and predictors of normal-weight central obesity– the AWI-Gen study findings. Glob Health Action.

2019;12. doi:10.1080/16549716.2019.1685809

- 31. Omar SM, Taha Z, Hassan AA, Al-Wutayd O, Adam I. Prevalence and factors associated with overweight and central obesity among adults in the Eastern Sudan. PLoS One. 2020;15: 1–10. doi:10.1371/journal.pone.0232624
- 32. Ahmed MH, Ali YA, Awadalla H, Elmadhoun WM, Noor SK, Almobarak AO. Prevalence and trends of obesity among adult Sudanese individuals: Population based study. Diabetes Metab Syndr Clin Res Rev. 2017;11: S963–S967. doi:10.1016/j.dsx.2017.07.023
- 33. Kabwama SN, Kirunda B, Mutungi G, Wesonga R, Bahendeka SK, Guwatudde D. Prevalence and correlates of abdominal obesity among adults in Uganda: Findings from a national cross-sectional, population based survey. BMC Obes. 2018;5: 1–9. doi:10.1186/s40608-018-0217-1
- 34. Chukwuonye II, Chuku A, Onyeonoro UU, Okpechi IG, Madukwe OO, Umeizudike TI, et al. Prevalence of abdominal obesity in Abia state, Nigeria: Results of a population-based house-to-house survey. Diabetes, Metab Syndr Obes Targets Ther. 2013;6: 285–291. doi:10.2147/DMSO.S43545
- 35. Gbary AR, Kpozehouen A, Houehanou YC, Djrolo F, Amoussou MP, Tchabi Y, et al. Prevalence and risk factors of overweight and obesity: findings from a cross-sectional community-based survey in Benin. Glob Epidemic Obes. 2014;2: 3. doi:10.7243/2052-5966-2-3
- Mekonnen T, Animaw W, Seyum Y. Overweight/obesity among adults in North-Western Ethiopia: A community-based cross sectional study. Arch Public Heal. 2018;76: 1–6. doi:10.1186/s13690-018-0262-8
- 37. Wong MCS, Huang J, Wang J, Chan PSF, Lok V, Chen X, et al. Global, regional and time-trend prevalence of central obesity: a systematic review and meta-analysis of 13.2 million subjects. Eur J Epidemiol. 2020;35: 673–683. doi:10.1007/s10654-020-00650-3
- 38. Ana M. Lo´pez-Sobaler, Elena Rodri´guez-Rodri´guez, Javier Aranceta-Bartrina, A´ ngel Gil, Marcela Gonza´lez-Gross, Lluis Serra-Majem, Gregorio Varela-Moreiras RMO. General and abdominal obesity is related to physical activity, smoking and sleeping behaviours and mediated by the educational level: Findings from the ANIBES study in Spain. PLoS One. 2016;11: 1–13. doi:10.1371/journal.pone.0169027
- 39. Pradeepa R, Anjana RM, Joshi SR, Bhansali A, Deepa M, Joshi PP. prevalence of generalized & abdominal obesity in urban & rural. 2015; 139–150. doi:10.4103/0971-5916.164234
- 40. Mogre V, Aleyira S, Nyaba R. Factors associated with central overweight and obesity in students attending the University for Development Studies in Tamale, Ghana: A cross-sectional study. South African J Clin Nutr. 2014;27: 69–74. doi:10.1080/16070658.2014.11734490
- 41. Wang H, Wang J, Liu MM, Wang D, Liu YQ, Zhao Y, et al. Epidemiology of general obesity, abdominal obesity and related risk factors in urban adults from 33 communities of northeast china: The CHPSNE study. BMC Public Health. 2012;12: 1. doi:10.1186/1471-2458-12-967
- 42. Bezerra IN, Curioni C, Sichieri R. Association between eating out of home and body weight. Nutrition Reviews. 2012. pp. 65–79. doi:10.1111/j.1753-4887.2011.00459.x
- 43. Pengpid S, Peltzer K. Associations between behavioural risk factors and overweight and obesity among adults in population-based samples from 31 countries. Obes Res Clin Pract. 2016; 1–9. doi:10.1016/j.orcp.2016.08.001
- 44. Clair C, Chiolero A, Faeh D, Cornuz J, Marques-Vidal P, Paccaud F, et al. Dose-dependent positive

association between cigarette smoking, abdominal obesity and body fat: Cross-sectional data from a population-based survey. BMC Public Health. 2011;11. doi:10.1186/1471-2458-11-23

- 45. Francisco José Gondim P, Cristiano Penas Seara P, Carmem Cristina B. Physical Activity in the Prevention of Abdominal Obesity: Type, Duration and Intensity. Int J Sport Exerc Med. 2018;4: 1–4. doi:10.23937/2469-5718/1510106
- 46. Kim K-B, Kim K, Kim C, Kang S-J, Kim HJ, Yoon S, et al. Effects of Exercise on the Body Composition and Lipid Profile of Individuals with Obesity: A Systematic Review and Meta-Analysis. J Obes Metab Syndr. 2019;28: 278–294. doi:10.7570/jomes.2019.28.4.278
- 47. Girma T, Mossie A, Getu Y. Association between body composition and khat chewing in Ethiopian adults. BMC Res Notes. 2015; 1–5. doi:10.1186/s13104-015-1601-2
- 48. Jalali-Farahani S, Amiri P, Karimi M, Gharibzadeh S, Mirmiran P, Azizi F. Socio-Behavioral Factors Associated with Overweight and Central Obesity in Tehranian Adults: a Structural Equation Model. Int J Behav Med. 2017;24: 110–119. doi:10.1007/s12529-016-9574-7
- 49. Bezerra IN, Curioni C, Sichieri R. Association between eating out of home and body weight. Nutrition Reviews. 2012. pp. 65–79. doi:10.1111/j.1753-4887.2011.00459.x
- 50. Kuźbicka K, Rachoń D. Bad eating habits as the main cause of obesity among children. Pediatr Endocrinol Diabetes Metab. 2013;19: 106–110.
- Sartorius B, Veerman LJ, Manyema M, Chola L, Hofman K. Determinants of obesity and associated population attributability, South Africa: Empirical evidence from a national panel survey, 2008-2012. PLoS One. 2015;10: 2008–2012. doi:10.1371/journal.pone.0130218
- 52. Obesity mapping : a conceptual framework. 2020.
- 53. Sullivan KM, Soe MM. Sample Size for a Cross-Sectional , Cohort , or Clinical Trial Studies Sample Size for Cross-Sectional & Cohort Studies & Clinical Trials. 2007.
- 54. Kumar R. Respondent Selection Methods in Household Surveys. Jharkhand Journal Dev Manag Stud. 2014;XII: 5701–5708. Available: https://www.researchgate.net/publication/267512058
- 55. WHO. Global Physical Activity Questionnaire (GPAQ) Analysis Guide. Geneva World Heal Organ. 2012.
- 56. WFP, FAO. Food consumption score: Construction of the FCS. World Food Program. 2008; 1–102. Available: http://documents.wfp.org/stellent/groups/public/documents/ena/wfp196627.pdf
- 57. Source: Lovibond, S.H., & Lovibond, P.F. (1995). Manual for the Depression Anxiety Stress Scales (2nd. ed). Sydney, Australia: Psychology Foundation.
- 58. Endalifer ML, Diress G. Mechanism of Obesity : A Systematic Review. J Obes. 2020;2020: 1–8.
- 59. Veghari G, Sedaghat M, Banihashem S, Moharloei P, Angizeh A, Tazik E, et al. Prevalence of general, abdominal and waist-to-hip ratio obesity among adults in the north of Iran: an ethnical comparative study. Int J Med Res Heal Sci. 2016;5: 219–225.
- 60. K. Malik S, Kouame J, Gbane M, Coulibaly M, D. Ake M, Ake O. Prevalence of abdominal obesity and its correlates among adults in a peri-urban population of West Africa. AIMS Public Heal. 2019;6: 334–344. doi:10.3934/publichealth.2019.3.334
- 61. Sunday U. Magnitude of Obesity, Abdominal Adiposity and their Association with Hypertension and

Diabetes- A Cross Sectional Study. J Metab Syndr. 2014;03. doi:10.4172/2167-0943.1000146

- 62. Undavalli VK, Ponnaganti SC, Narni H. Prevalence of generalized and abdominal obesity: India's big problem. Int J Community Med Public Heal. 2018;5: 1311–1316. doi:10.18203/2394-6040.ijcmph20180984
- 63. Kabwama SN, Kirunda B, Mutungi G, Wesonga R, Bahendeka SK, Guwatudde D. Prevalence and correlates of abdominal obesity among adults in Uganda: findings from a national cross-sectional, population based survey 2014. BMC Obes. 2018;5: 40. doi:10.1186/s40608-018-0217-1
- 64. Prasad DS, Kabir Z, Revathi Devi K, Peter PS, Das BC. Gender differences in central obesity: Implications for cardiometabolic health in South Asians. Indian Heart J. 2020;72: 202–204. doi:10.1016/j.ihj.2020.04.008
- 65. Csongová M, Volkovová K, Gajdoš M, Gurecká R, Koborová I, Líšková A, et al. Gender-associated differences in the prevalence of central obesity using waist circumference and waist-to-height ratio, and that of general obesity, in Slovak adults. Cent Eur J Public Health. 2018;26: 228–233. doi:10.21101/cejph.a4719
- 66. Peters SAE, Bots SH, Woodward M. Sex differences in the association between measures of general and central adiposity and the risk of myocardial infarction: Results from the UK Biobank. J Am Heart Assoc. 2018;7: 6–10. doi:10.1161/JAHA.117.008507
- 67. Villareal DT, Apovian CM, Kushner RF, Klein S. Obesity in older adults: Technical review and position statement of the American Society for Nutrition and NAASO, the Obesity Society. Obes Res. 2005;13: 1849–1863. doi:10.1038/oby.2005.228
- 68. Owolabi EO, Ter Goon D, Adeniyi OV. Central obesity and normal-weight central obesity among adults attending healthcare facilities in Buffalo City Metropolitan Municipality, South Africa: A cross-sectional study. J Heal Popul Nutr. 2017;36: 1–10. doi:10.1186/s41043-017-0133-x
- 69. Chukwuonye II, Chuku A, John C, Ohagwu KA, Imoh ME, Isa SE, et al. Prevalence of overweight and obesity in adult Nigerians A systematic review. Diabetes, Metab Syndr Obes Targets Ther. 2013;6: 43–47. doi:10.2147/dmso.s38626
- 70. Gunderson EP. Childbearing and Obesity in Women: Weight Before, During, and After Pregnancy. Obstet Gynecol Clin North Am. 2009;36: 317–332. doi:10.1016/j.ogc.2009.04.001
- 71. Jakicic JM, Otto AD. Physical activity considerations for the treatment and prevention of obesity. Am J Clin Nutr. 2005;82: 226–229. doi:10.1093/ajcn.82.1.226s
- 72. Kanter R, Caballero B. Global Gender Disparities in Obesity : A Review. Am Soc Nutr. 2012;3: 491–498. doi:10.3945/an.112.002063.published
- 73. Mathieu P, Lemieux I, Després JP. Obesity, inflammation, and cardiovascular risk. Clin Pharmacol Ther. 2010;87: 407–416. doi:10.1038/clpt.2009.311
- 74. Jura M, Kozak LP. Obesity and related consequences to ageing. Age (Omaha). 2016;38. doi:10.1007/s11357-016-9884-3
- 75. Anderson CAM, Mongraw-Chaffin M. Central obesity in older adults: What should be the priority? Journal of the American Heart Association. 2018. pp. 1–3. doi:10.1161/JAHA.118.010119
- 76. Mulugeta A, Zhou A, Power C, Hyppönen E. Obesity and depressive symptoms in mid-life: A population-based cohort study. BMC Psychiatry. 2018;18: 1–10. doi:10.1186/s12888-018-1877-6

- 77. Hryhorczuk C, Sharma S, Fulton SE. Metabolic disturbances connecting obesity and depression. Front Neurosci. 2013;7: 1–14. doi:10.3389/fnins.2013.00177
- 78. Review AS. Overweight, Obesity, and Depression. 2010;67: 220–229.
- Bornstein SR, Schuppenies A, Wong ML, Licinio J. Approaching the shared biology of obesity and depression: The stress axis as the locus of gene-environment interactions. Molecular Psychiatry. 2006. pp. 892–902. doi:10.1038/sj.mp.4001873
- 80. Mogre V, Oladele J, Amalba A. Impact of physical activity levels and diet on central obesity among civil servants in Tamale metropolis. J Med Biomed Sci. 2012;1: 1–9.
- 81. Ailshul MD. Energy balance and obesity in man. Nutr Today. 1975;10: 29–30. doi:10.1097/00017285-197509000-00009
- 82. Cox CE. Role of physical activity for weight loss and weight maintenance. Diabetes Spectr. 2017;30: 157–160. doi:10.2337/ds17-0013

#### 11. Annex

Hello! My name is \_\_\_\_\_\_ and I am collecting data for a study on the <u>central</u> obesity and its associated factors among adults aged 18-64 years in Debre Tabor Town, northeast, Ethiopia, 2021. I would like to ask you some questions about the risk factors of Central obesity and if you are willing, we will also take some physical examination. The interview would take about 15 minutes. Your participation in the study is very critical to know the magnitude of Central Obesity and its associated factors. However, your participation is fully voluntary. The data you provided will be kept in a highly confidential manner and none of your identifiers will be on the questionnaire.

Verbal consent obtained		Yes	No
Date of data collection	//		

|--|

Part I: Socio-demographic characteristics					
A0	Unique Identification Number	ID			
A1	What is your Sex				
A2	What is your Ethnic group	2. Female1. Amhara2. Oromo3. Tigre4. Others (specify)			
A3	What is your Age	years			
A4	What is your Religion	<ol> <li>Orthodox</li> <li>Protestant</li> <li>Moslim</li> <li>Catholic</li> <li>Other specify</li> </ol>			
A5	What is your Occupation	<ol> <li>Employ (both government and non governmental)</li> <li>Merchant</li> <li>Daily labourer</li> <li>Housewife</li> <li>Student</li> <li>Others(specify)</li> </ol>			
A6	What is your Marital status	1. Married			

		2. Single
		3 Widowed
		4 Divorced
		5 Separated
A8	What is your Level of educational status	1 Can't read and write
110	what is your Level of eddealional status	2 can read and write but no
		formal education
		3 primary(1-8)
		4  Secondary(9.12)
		5 Tertiary and above
	Part II· Rehavioral Measurements	
	Tobacco, chewing Khat and Alcohol use practice	
B1	Have you ever smoked any tobacco products such as	0 No
	cigarettes cigars shisha or nines?	1  Yes
B2	Do you currently smoke cigarettes?	$\frac{1.105}{0.00}$
D2	Do you currently smoke ergurenes.	1  Yes
B3	If yes to B2 do you currently smoke tobacco products	$\frac{1.105}{0.No}$
	daily?	1  Yes
R4	For how long do you smoke cigarettes	$\frac{1.105}{0.No}$
	Tor now rong do you smoke ergurenes	1  Yes
B5	Do you chew Khat?	$\frac{1.105}{0.00}$
<b>D</b> 5		1  Yes
B6	If yes, in a typical week how many days do you chew	days
20	Khat?	
B7	How many grams do you chew on one of those days?	grams
B8	Have you consumed any alcohol within the past 12	0. No
	months?	1. Yes
B9	During the past 12 months, have you had a standard	0. No
	alcoholic drink? (Vodka, Whisky, Araki ETC.)	1. Yes
	Part III: food consumption status	
D1	In a typical week, on how many days do you eat fruit?	Days
D2	How many servings of fruit do you eat on one of those days?	Servings
D3	In a typical week, on how many days do you eat	
	vegetables?	Days
		2
D4	How many servings of vegetables do you get on one of	Carving
D4	those days?	
D5	In a typical week on how many days do you eat	Davs
D5	protein source foods from animals (beef lamb	Days
	chicken fish and egg)?	
D6	In a typical week on how many days do you eat	Davs
D0	protein source foods from plants (pea bean chicknea	Days
	nuts groundnuts)?	
D7	In a typical week on how many days do you eat	Dave
	energy source foods (cereal grains energy-rich tubers	Days
	such as potatoes, sweet potatoes)?	
	saen as polatoes, sweet polatoes).	

D8	In a typical week, on how many days do you eat milk and milk products (milk cheese, Yogurt)?	Days
D9	In a typical week on how many days do you eat fats	Davs
27	(fats and oils)?	Duy5
D10	In a typical week, on how many days do you eat /drink	Days
	discretionary calories (soft drinks, sugar, chocolates,	
	honey)?	
D11	In a typical week, on how many days do you eat foods	Days
	fried /baked in an oil (egg? Chips, Biscuits, crackers,	5
	cakes)?	
Oil Us	e Practice	
D12	What type of oil or fat is often used for meal	1. Liquid Vegetable oil
	preparation in your household?	2. Solidified vegetable oil
		3. Butter
		4. Margarine / peanut
		butter
		5. Sheno lega
		6. Other
		7. None in particular
		8. None used
D10		9. Don't know
D13	On average, how many meals per week do you eat that	Meals
	were not prepared at a nome?	
Part I	V: Physical Activity	0 N
PI	Does your work involve a vigorous-intensity activity	0. No
	that causes large increases in breathing or heart rate	1. Yes
	like [carrying or lifting heavy loads, digging or	
	construction work for at least 10 minutes	
D2	continuously?	Dava
P2	in a typical week, on now many days do you do	Days
D2	How much time do you spand doing vigorous intensity	Hours Minutos
F 5	activities at work on a typical day?	IIOursMinutes
P4	Does your work involve a moderate-intensity activity	0 No
1 1	that causes small increases in breathing or heart rate	1. Yes
	such as brisk walking <i>[or carrying light loads]</i> for at	
	least 10 minutes continuously?	
P5	In a typical week, on how many days do you do	Days
	moderate-intensity activities as part of your work?	
P6	How much time do you spend doing moderate-	HoursMinutes
	intensity activities at work on a typical day?	
P7	Do you walk or use a bicycle (pedal cycle) for at least	0. No
	10 minutes continuously to get to and from places?	1. Yes
P8	In a typical week, on how many days do you walk or	Days
	bicycle for at least 10 minutes continuously to get to	
	and from places?	
P9	How much time do you spend walking or bicycling for	HoursMinutes

	travel on a typical day?					
P10	Do you do any vigorous-intensity sports, fitness or	0.	No			
	recreational (leisure) activities that cause large 1					
	increases in breathing or heart rate like [running or					
	<i>football]</i> for at least 10 minutes continuously?					
P11	In a typical week, on how many days do you do			Da	ays	
	vigorous-intensity sports, fitness or recreational					
<b>D10</b>	(leisure) activities?					
P12	How much time do you spend doing vigorous-intensity	H	ours	_Minut	es	
	sports, fitness or recreational activities on a typical					
D12	Do you do any moderate intensity sports fitness or	0	No			
F13	recreational <i>(laisura)</i> activities that cause a small	0. 1	NO			
	increase in breathing or heart rate such as brisk	1.	105			
	walking <i>[cycling swimming and yolleyball]</i> for at					
	least 10 minutes continuously?					
P14	In a typical week, on how many days do you do			Da	ays	
	moderate-intensity sports, fitness or recreational				5	
	(leisure) activities?					
P15	How much time do you spend doing moderate-	H	ours	_Minut	es	
	intensity sports, fitness or recreational (leisure)					
	activities on a typical day?					
Seden	tary Behavior					
S1	How much time do you usually spend sitting or	H	ours	_Minut	es	
	reclining on a typical day?					
<u>S2</u>	How many hours do usually sleep in a typical day			_Hours		
<b>S</b> 3	Do you often have a fragmented sleep with multiple 1. N					
	wake ups and falling asleep cycles? 2. So					
<b>C</b> 4	How many days due you have right duties in one	3.Yes	most of	the tin	ne	
54	How many days due you have hight duties in one month(only for hospital workers)			-days		
Denres	sion Anxiety and Stress Scale - 21 Items (DASS-21)					
Plea	use read each statement and circle a number 0, 1, 2 or 3	3 whic	n indic	ates ho	w muc	h the
stat	ement applied to you over the past week.		i illuio		iiide.	
The	rating scale is as follows:					
	id not apply to me at all					
	not upply to file at an					
	pplied to me to a considerable degree or a good next of time					
	pplied to me to a considerable degree of a good part of tim	le				
3 A	pplied to me very much or most of the time				•	
1 (s)	I found it hard to wind down		0	1	2	3
2 (a)	I was aware of dryness of my mouth		0	1	2	3
3 (d)	I couldn't seem to experience any positive feeling at all		0	1	2	3
<b>4</b> (a)	I experienced breathing difficulty (e.g., excessively	rapid	0	1	2	3
	breathing, breathlessness in the absence of phy	ysical				
	exertion)		0	1	•	
5 (d)	I found it difficult to work up the initiative to do things		0	1	2	3

•					
2	3				
2	3				
2	3				
2	3				
2	3				
2	3				
2	3				
2	3				
2	3				
2	3				
2	3				
2	3				
2	3				
2	3				
2	3				
Household wealth, Does the household have any of the following properties?					
<u> </u>	es				
$\frac{1}{1}$	es				
$\frac{1}{1}$	es				
$\frac{1}{1}$	es				
$\frac{1.1}{1.1}$	es es				
$\frac{1}{1}$ $\frac{1}{1}$ $\frac{1}{1}$	es l				
$\frac{1.1}{1.1}$	es les				
$\frac{1}{1}$ 1. Y	Tes Tes				
D 1. Y	es				
D 1. Y	es				
D 1. Y	es				
D 1. Y	es				
D 1. Y	es				
<u> </u>	fes				
$\frac{1.Y}{1.Y}$	es				
1.Y	es				
$\frac{1}{2}$	es				
$\frac{1.1}{1.1}$	es es				
, I. I	63				
cm					
cm					
	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2				

-	-

#### 12. Amharic version questionnaire

ሰላም ነው! ስሜ \_\_\_\_\_\_ እባላለው፡፡ እኔ ከመጠን በላይ ውፍረት እና የአካል ብቃት እንቅስቃሴ ደረጃ ንፅፅር እና ከ 18-64 ዓመት ዕድሜ ባላቸው ጎልማሳዎች መካከል በሰሜን ምስራቅ ፣ ኢትዮጵያ በማወዳደር ጥናት ላይ ለመስራት ነው ፡፡ ስለ ማዕከላዊ ውፍረት ተጋላጭነት ምክንያቶች አንዳንድ ጥያቄዎችን ልጠይቅዎ ከፈለጉ እና ፈቃደኛ ከሆኑ እኛ ደግሞ የአካል ምርመራ እናደርጋለን። ቃለ መጠይቁ 15 ደቂቃ ያህል ይወስዳል ፡፡ የማዕከላዊ ውፍረት እና ተጓዳኝ ምክንያቶቹን ለማወቅ በጥናቱ ውስጥ የእርስዎ ተሳትፎ በጣም ወሳኝ ነው ፡፡ ሆኖም የእርስዎ ተሳትፎ ሙሉ በሙሉ በፈቃደኝነት የሚደረግ ነው። ያቀረቡት መረጃ በከፍተኛ በሚስጥራዊ ሁኔታ ይቀመጣል እና ከግል መለያዎ ውስጥ አንዳቸውም በመጠይቁ ላይ አይኖሩም።

የቃል ስምምነት የመረጃ መሰብሰቢያ ቀን \_\_\_\_\_ / \_\_\_\_\_ / \_\_\_\_\_ አዎ የለም (በኢትዮጵያ አቆጣጠር የቀን ወር ዓመት)

ከፍል	አንድ፡ አ <b>ጢቃላይ <i>መ</i>ረ</b> ጃ	
A0	መለያ ቁጥር	ቁጥር
A1	የታ	1. ወንድ 2. ሴት
A2	ብሄር	1. አማራ 2. ኦሮሞ 3. <i>ትግፌ</i> 4. ሌላ ካለ ይጠቅስ
A3	እድሜ	በአመት
A4	ሃይማኖት	1. አርቶዶክስ 2. ፕሮቴስታንት 3. እስላም 4. ካቶሊክ 5. ሌላ ካለ ይጠቅስ
A5	ስራ-	<ol> <li>ተቀጣሪ (የመንግስት / መንግስታዊ ያልሆነ)</li> <li>ነጋዴ</li> <li>የቀን ሰራተኛ</li> <li>የቤት እመቤት</li> <li>ተማሪ</li> <li>ሌላ ካለ ይጠቀስ</li> </ol>
A6	የትዳር ሁኔታ	1. ያገባ 2. ያላገባ 3. ጋለምታ 4. የተፋታ 5. ተለያይተው የሚኖሩ
A7	አማካኝ የአመት <i>ገ</i> ቢ	በብር
A8	የትምህርት ደረጃ	1. ማንበብና መጻፍ የማይችል

		2. ማንበብና መጻፍ የሚችል
		3 የመኛመሪያ ደረጃ
		4 <i>ሁለተኛ ደረጃ</i>
		5 ለስተኛ ደረጃ እና በላይ
ከፍል	ሁለት፡ የባህርይ <i>መ</i> ለኪያዎች	
B1	እንደ ሲ <i>ጋ</i> ራ ፤ ሺሻ ወይም ሌሎች ማናቸውንም የትምባሆ ምርቶች	1. በፋጹም
	አጭሰው ያውቃ።	2. አዎ
B2	አሁን ሲጋራ ያጨሳሉ?	1. በፋጹም
		2. አዎ
B3	ለ B2 መለስዎ አዎ ከሆነ, በቀን በቀን ያጨሳሉ?	1. በፋጹም
		2. አዎ
B4	ለምን ያክል ጊዜ ሲ <i>ጋራ አ</i> ጭሰው ያውቃሉ፡፡	1. በፋጹም
		2. አዎ
B5	ሜት ይቅጣሉ?	2. በፋጹም
		3. አዎ
B6	እልሲቆ አዎ ከሆነ፣ በሳምንት ለምል ያከል ጊዜ ይጠቀጣሉ?	በቀን
B7	ምን ያክል ባራም አጭሰው ያውቃሉ?	በግራም
B8	ባለፉት 12 ወራት ውስፕ ማንኛውንም አልኮል ጠፕተዋል?	1. በፋጹም
		2. አዎ
B9	ለባፉት 12 ወራት ውስጥ መደበኛ የአልኮሆል መጠጥ አለዎት? (ቮድካ	1. በፋጹም
	፤ ውስኪ ፤ አረኪ ወዘተ)	2. አዎ
	,	
	ክፍል ለኢት・ የአመንግብ ኪነ ታን የመመልክታ መጥደ ቅች።	
<del>ہ</del> /ھ	<u>ክፍል ሶስት፡ የአመጋገብ ሁኔታን የሚመለከቱ መጠይቅች፡፡</u> መመደቅ	መለሽ/ አማረጭ
ተ/ቁ	<u>ክፍል ሶስት፡ የአመጋገብ ሁኔታን የሚመለከቱ መጠይቅች፡፡</u> መጠይቅ	ምላሽ/ አማራጭ
<b>ተ/ቁ</b> 1	<u>ክፍል ሶስት፡ የአመጋገብ ሁኔታን የሚመለከቱ መጠይቅች፡፡</u> መጠይቅ በቀን ስንት ጊዜ ትመገቢያልሽ1	<b>ምላሽ/ አጣራጭ</b> 0. አንድ 1. ሁለት ጊዜ 3. ሶስትና በላይ
<b>     ተ/ቁ</b> 1     D2	<u>ክፍል ሶስት፡ የአመጋገብ ሁኔታን የሚመለከቱ መጠይቅች፡፡</u> መጠይቅ በቀን ስንት ጊዜ ትመገቢያልሽ1 በአዘቦት ሳምንት፣ ለምን ያክል ቀን ፍራፍሬ ትመገቢያለሽ	<b>ምላሽ/ አጣራጭ</b> ዐ. አንድ 1. ሁለት ጊዜ 3. ሶስትና በላይ 
<b>↑/⊉</b> 1 D2 D3	<u>ክፍል ሶስት፡ የአመጋገብ ሁኔታን የሚመለከቱ መጠይቅች፡፡</u> መጠይቅ በቀን ስንት ጊዜ ትመገቢያልሽ1 በአዘቦት ሳምንት፣ ለምን ያክል ቀን ፍራፍሬ ትመገቢያለሽ ፍራፍሬ በምትመገቢበት ቀናት፣ ምን ያክል ትመገቢያለሽ (መጠን)	<b>ምላሽ/ አጣራጭ</b> 0. አንድ 1. ሁለት ጊዜ 3. ሶስትና በላይ   ቁራሽ /በፍሬ
<b>↑/‡</b> 1 D2 D3	<u>ክፍል ሶስት፡ የአመጋገብ ሁኔታን የሚመለከቱ መጠይቅች፡፡</u> መጠይቅ በቀን ስንት ጊዜ ትመገቢያልሽ1 በአዘቦት ሳምንት፣ ለምን ያክል ቀን ፍራፍሬ ትመገቢያለሽ ፍራፍሬ በምትመገቢበት ቀናት፣ ምን ያክል ትመገቢያለሽ (መጠን)	<b>ምላሽ/ አማራጭ</b> ዐ. አንድ 1. ሁለት ጊዜ 3. ሶስትና በላይ ቀናት ቁራሽ /በፍሬ
<b>⁺/</b> ₽ 1 D2 D3	<u>ክፍል ሶስት፡ የአመጋገብ ሁኔታን የሚመለከቱ መጠይቅች፡፡</u> መጠይቅ በቀን ስንት ጊዜ ትመገቢያልሽ1 በአዘቦት ሳምንት፣ ለምን ያክል ቀን ፍራፍሬ ትመገቢያለሽ ፍራፍሬ በምትመገቢበት ቀናት፣ ምን ያክል ትመገቢያለሽ (መጠን)	<b>ምላሽ/ አጣራጭ</b> 0. አንድ 1. ሁለት ጊዜ 3. ሶስትና በላይ ሰቀናት ቁራሽ /በፍሬ
<b>↑</b> / <b>₽</b> 1 D2 D3	<u>ክፍል ሶስት፡ የአመጋገብ ሁኔታን የሚመለከቱ መጠይቅች፡፡</u> መጠይቅ በቀን ስንት ጊዜ ትመገቢያልሽ1 በአዘቦት ሳምንት፣ ለምን ያክል ቀን ፍራፍሬ ትመገቢያለሽ ፍራፍሬ በምትመገቢበት ቀናት፣ ምን ያክል ትመገቢያለሽ (መጠን) በአዘቦት ሳምንት፣ ለምን የክል ቀን አትክልት ትመገቢያለሽ	<b>ምላሽ/ አጣራጭ</b> 0. አንድ 1. ሁለት ጊዜ 3. ሶስትና በላይ ቀናት ቁራሽ /በፍሬ 
<b>⁺/</b> ₽ 1 D2 D3 D4	<mark>ክፍል ሶስት፡ የአመጋገብ ሁኔታን የሚመለከቱ መጠይቅች፡፡</mark> መጠይቅ በቀን ስንት ጊዜ ትመገቢያልሽ1 በአዘቦት ሳምንት፣ ለምን ያክል ቀን ፍራፍሬ ትመገቢያለሽ ፍራፍሬ በምትመገቢበት ቀናት፣ ምን ያክል ትመገቢያለሽ (መጠን) በአዘቦት ሳምንት፣ ለምን ያክል ቀን አትክልት ትመገቢያለሽ	<b>ምላሽ/ አማራጭ</b> 0. አንድ 1. ሁለት ጊዜ 3. ሶስትና በላይ
ተ/⋬ 1 D2 D3 D4 D5	<u>ክፍል ሶስት፡ የአመጋገብ ሁኔታን የሚመለከቱ መጠይቅች፡፡</u> መጠይቅ በቀን ስንት ጊዜ ትመገቢያልሽ1 በአዘቦት ሳምንት፣ ለምን ያክል ቀን ፍራፍሬ ትመገቢያለሽ ፍራፍሬ በምትመገቢበት ቀናት፣ ምን ያክል ትመገቢያለሽ (መጠን) በአዘቦት ሳምንት፣ ለምን ያክል ቀን አትክልት ትመገቢያለሽ አትክልት በምትመገቢበት ቀናት፣ ምን ያክል ትመገቢያለሽ (መጠን)	<b>ምላሽ/ አጣራጭ</b> 0. አንድ 1. ሁለት ጊዜ 3. ሶስትና በላይ
<ul> <li><i>↑</i>/<i>₱</i></li> <li>1</li> <li>D2</li> <li>D3</li> <li>D4</li> <li>D5</li> <li>D6</li> </ul>	<u>ክፍል ሶስት፡ የአመጋገብ ሁኔታን የሚመለከቱ መጠይቅች፡፡</u> መጠይቅ በቀን ስንት ጊዜ ትመገቢያልሽ1 በአዘቦት ሳምንት፣ ለምን ያክል ቀን ፍራፍሬ ትመገቢያለሽ ፍራፍሬ በምትመገቢበት ቀናት፣ ምን ያክል ትመገቢያለሽ (መጠን) በአዘቦት ሳምንት፣ ለምን ያክል ቀን አትክልት ትመገቢያለሽ (መጠን) በአዘቦት ሳምንት፣ ለምን ያክል ቀን የእንሰሳት ተዋጽኦ ፕሮቲን (ሲጋ፣	<b>ምላሽ/ አማራጭ</b> 0. አንድ 1. ሁለት ጊዜ 3. ሶስትና በላይ
r//#         1         D2         D3         D4         D5         D6	<u>ክፍል ሶስት፡ የአመጋገብ ሁኔታን የሚመለከቱ መጠይቅች፡፡</u> መጠይቅ በቀን ስንት ጊዜ ትመንቢያልሽ1 በአዘቦት ሳምንት፣ ለምን ያክል ቀን ፍራፍሬ ትመንቢያለሽ ፍራፍሬ በምትመንቢበት ቀናት፣ ምን ያክል ትመንቢያለሽ (መጠን) በአዘቦት ሳምንት፣ ለምን ያክል ቀን አትክልት ትመንቢያለሽ (መጠን) በአዘቦት ሳምንት፣ ለምን ያክል ቀን የእንስሳት ተዋጽኦ ፕሮቲን (ሲጋ፣ መታት፣ እንቀለላ፣ አስ መዘታ) ታመንቦ ይላሽ	<b>ምላሽ/ አማራጭ</b> 0. አንድ 1. ሁለት ጊዜ 3. ሶስትና በላይ
<ul> <li><i>↑</i>/<i>₱</i></li> <li>1</li> <li>D2</li> <li>D3</li> <li>D4</li> <li>D5</li> <li>D6</li> </ul>	ከፍል ሶስት: የአመጋገብ ሁኔታን የሚመለከቱ መጠይቅች::           መጠይቅ           በቀን ስንት ጊዜ ትመገቢያልሽ1           በአዘቡት ሳምንት፣ ለምን ያክል ቀን ፍራፍሬ ትመገቢያለሽ           ፍራፍሬ በምትመገቢበት ቀናት፣ ምን ያክል ትመገቢያለሽ (መጠን)           በአዘቡት ሳምንት፣ ለምን ያክል ቀን አትክልት ትመገቢያለሽ (መጠን)           በአዘቡት ሳምንት፣ ለምን ያክል ቀን አትክልት ትመገቢያለሽ (መጠን)           በአዘቡት ሳምንት፣ ለምን ያክል ቀን አትክልት ትመገቢያለሽ (መጠን)           በአዘቡት ሳምንት፣ ለምን ያክል ቀን የእንስሳት ተዋጽኦ ፕሮቲን (ሲጋ፣ መተት፣ እንቁላል፣ አሳ ወዘተ) ትመገቢያለሽ	<b>ምላሽ/ አማራጭ</b> 0. አንድ 1. ሁለት ጊዜ 3. ሶስትና በላይ
ተ/ቁ         1         D2         D3         D4         D5         D6         D7	ከፍል ሶስት: የአመጋገብ ሁኔታን የሚመለከቱ መጠይቅች::         መጠይቅ         በቀን ስንት ጊዜ ትመገቢያልሽ1         በአዘቦት ሳምንት፣ ለምን ያክል ቀን ፍራፍሬ ትመገቢያለሽ         ፍራፍሬ በምትመገቢበት ቀናት፣ ምን ያክል ትመገቢያለሽ (መጠን)         በአዘቦት ሳምንት፣ ለምን ያክል ቀን አትክልት ትመገቢያለሽ (መጠን)         በአዘቦት ሳምንት፣ ለምን ያክል ቀን አትክልት ትመገቢያለሽ (መጠን)         በአዘቦት ሳምንት፣ ለምን ያክል ቀን የእንሰሳት ተሞጽኦ ፕሮቲን (ሲጋ፣ ወተት፣ እንቁላል፣ አሳ ወዘተ) ትመገቢያለሽ         በአዘቦት ሳምንት፣ ለምን ያክል ቀን የእንሰሳት ተሞጽኦ ፕሮቲን (ባቋላ፣	<b>ምላሽ/ አማራጭ</b> 0. አንድ 1. ሁለት ጊዜ 3. ሶስትና በላይ
<ul> <li><i>↑</i>/<i>‡</i></li> <li>1</li> <li>D2</li> <li>D3</li> <li>D4</li> <li>D5</li> <li>D6</li> <li>D7</li> </ul>	ክፍል ሶስት: የአመጋገብ ሁኔታን የሚመለከቱ መጠይቅች::         መጠይቅ         በቀን ስንት ጊዜ ትመንቢያልሽ1         በአዘቦት ሳምንት፣ ለምን ያክል ቀን ፍራፍሬ ትመንቢያለሽ         ፍራፍሬ በምትመንቢበት ቀናት፣ ምን ያክል ትመንቢያለሽ (መጠን)         በአዘቦት ሳምንት፣ ለምን ያክል ቀን አትክልት ትመንቢያለሽ (መጠን)         በአዘቦት ሳምንት፣ ለምን ያክል ቀን የእንስሳት ተዋጽኦ ፕሮቲን (ሲጋ፣ ወተት፣ እንቁላል፣ አሳ ወዘተ) ትመንቢያለሽ         በአዘቦት ሳምንት፣ ለምን ያክል ቀን የእጽዋት ተዋጽኦ ፕሮቲን (ባቄላ፣ ኦተር፣ ሽመብረ፣ ኦቾላኑ መዘተ) ትመንባ የላሽ	<b>ምላሽ/ አማራጭ</b> 0. አንድ 1. ሁለት ጊዜ 3. ሶስትና በላይ 
<ul> <li><i>↑</i>/<i>₱</i></li> <li>1</li> <li>D2</li> <li>D3</li> <li>D4</li> <li>D5</li> <li>D6</li> <li>D7</li> </ul>	ከፍል ሶስት፡ የአመጋገብ ሁኔታን የሚመለከቱ መጠይቅች፡፡         መጠይቅ         በቀን ስንት ጊዜ ትመገቢያልሽ1         በአዘቡት ሳምንት፣ ለምን ያክል ቀን ፍራፍሬ ትመገቢያለሽ         ፍራፍሬ በምትመገቢበት ቀናት፣ ምን ያክል ትመገቢያለሽ (መጠን)         በአዘቡት ሳምንት፣ ለምን ያክል ቀን አትክልት ትመገቢያለሽ (መጠን)         በአዘቡት ሳምንት፣ ለምን ያክል ቀን የእንስሳት ተዋጽኦ ፕሮቲን (ሲጋ፣ ወተት፣ እንቁላል፣ አሳ ወዘተ) ትመገቢያለሽ         በአዘቡት ሳምንት፣ ለምን ያክል ቀን የእጽዋት ተዋጽኦ ፕሮቲን (ባቄላ፣ አተር፣ ሽምብራ፣ ኦቾለኒ ወዘተ) ትመገቢያለሽ	<b>ምላሽ/ አማራጭ</b> 0. አንድ 1. ሁለት ጊዜ 3. ሶስትና በላይ
<ul> <li><i>t</i>/<i>f</i></li> <li>1</li> <li>D2</li> <li>D3</li> <li>D4</li> <li>D5</li> <li>D6</li> <li>D7</li> <li>D8</li> </ul>	<u>ከፍል ሶስት፡ የአመጋንብ ሁኔታን የሚመለከቱ መጠይቅች፡፡</u> መጠይቅ በቀን ስንት ጊዜ ትመንቢያልሽ1 በአዘቦት ሳምንት፣ ለምን ያክል ቀን ፍራፍሬ ትመንቢያለሽ ፍራፍሬ በምትመንቢበት ቀናት፣ ምን ያክል ትመንቢያለሽ (መጠን) በአዘቦት ሳምንት፣ ለምን ያክል ቀን አትክልት ትመንቢያለሽ (መጠን) በአዘቦት ሳምንት፣ ለምን ያክል ቀን የእንስሳት ተዋጽኦ ፕሮቲን (ሲጋ፣ ወተት፣ እንቁላል፣ አሳ ወዘተ) ትመንቢያለሽ በአዘቦት ሳምንት፣ ለምን ያክል ቀን የእጽዋት ተዋጽኦ ፕሮቲን (ባቄላ፣ አተር፣ ሽምብራ፣ ኦቾለኒ ወዘተ) ትመንቢያለሽ	<b>ምላሽ/ አማራጭ</b> 0. አንድ 1. ሁለት ጊዜ 3. ሶስትና በላይ 
<ul> <li><i>⁺</i>/<i>‡</i></li> <li>1</li> <li>D2</li> <li>D3</li> <li>D4</li> <li>D5</li> <li>D6</li> <li>D7</li> <li>D8</li> </ul>	<u>ከፍል ሶስት፡ የአመጋንብ ሁኔታን የሚመለከቱ መጠይቅች፡፡</u> መጠይቅ በቀን ስንት ጊዜ ትመንቢያልሽ1 በአዘቦት ሳምንት፣ ለምን ያክል ቀን ፍራፍሬ ትመንቢያለሽ ፍራፍሬ በምትመንቢበት ቀናት፣ ምን ያክል ትመንቢያለሽ (መጠን) በአዘቦት ሳምንት፣ ለምን ያክል ቀን አትክልት ትመንቢያለሽ (መጠን) በአዘቦት ሳምንት፣ ለምን ያክል ቀን የእንስሳት ተዋጽኦ ፕሮቲን (ሲጋ፣ ወተት፣ እንቁላል፣ አሳ ወዘተ) ትመንቢያለሽ በአዘቦት ሳምንት፣ ለምን ያክል ቀን የእጽዋት ተዋጽኦ ፕሮቲን (ባቃላ፣ አተር፣ ሽምብራ፣ ኦቾለኒ ወዘተ) ትመንቢያለሽ	<b>ምላሽ/ አማራጭ</b> 0. አንድ 1. ሁለት ጊዜ 3. ሶስትና በላይ 

D9	በአዘቦት ሳምንት፣ ለምን ያክል ቀን ወተት እና የወተት ተዋጽኦ <i>ትመ</i> ገባለሽ		በቀናት	
<b>D</b> 10			0401	
D10	በአዘቦተ ሳምንተ፣ ለምን ያክል ቀን ቀባተነተ ያላቸው (ስብና ዘይተ) <i>ትመገ</i> ቢያለሽ		<u> </u>	
D11	በአዘቦት ሳምንት፣ ለምን ያክል ቀን ታፋጭነት ያለቸዉ ምግቦች አና		በቀናት	
	መጠጦች (ለስላሳ መጠጥ፣ ስኳር፣ ቸኮሌት፣ ማር ወዘተ) ትመንቢያሽ			
D12	በአዘቦት ሳምንት፣ ለምን ያክል ቀን በዘይት የተጠበሱ ምግቦችን		በቀናት	
	(ቸፕስ፣ ሳቡሳ፣ ወዘተ) ትመንቢያሽ			
የዘይት	<sup>.</sup> አጠ <i>ቃቀ</i> ም ልምድ			
D12	በቤተሰብዎ ውስጥ ለምግብ ዝግጅት ብዙ ጊዜ ጥቅም ላይ የሚውለው	1. ሳሽ የአ	ትክልት ዘይት	
	ምን ዓይነት ዘይት ወይም ስብ ነው?	2. የተጠና	ከረ የአትክልት ዘይት	
		3. Plb	<u>ዓ / ወኔ ጆሔ</u> ቴ ሕበ	
		4. ግቢታሪ 5 ሸኖለኋ	) )	
		6. AA		
		ነም አንጠቅመም		
D13	በአማካይ በሳምንት ስንት ምግብ በቤት ውስጥ ያልተዘጋጁ ምግብ	ምግቦች		
	ይመገባሉ?			
ክፍል	አራት: የአካል ብቃት እንቅስቃሴ			
P1	ሥራዎ እንደ ትንፋሽ ወይም የልብ ምትን የመሰለ ከፍተኛ ጭንቀትን [	ከባድ ሸክምችን	2. አያካትትም	
	መሸከም ወይም ማንሳት ፣ መቆፈር ወይም የግንባታ ሥራ] ቢያንስ ለ 10 ደቂቃዎች		3. ያካትታል	
Da	ያለማዋረጥ የሚጨምር ኃይለና እንቀስቃሴን ያካትታል?   በተለመደሙ ለመንጉ ሙስም መረጣ ተባላጅ የሆኑሉ በ በተኑ እንትእ ተላጥ	ኒቲካ አካኒ ሐርኒ		
P2	በተለመዱው ባምንተ ውጤት ሥራዎ ንይበት የለባል ዝጅተ ለንዋቢይቤዎ ዕъህ የኢን	ጥ 7 (በ 7ጥ ዋዓጥ	(l ሐርት	
P3	ጋ የፈጣበ። በተለመየሙ ቀን በሥራ ለይ ኃይለኛ እንቅስቀሲዎችን ለማድረማ ዓ	ምን የህል ገዚ	ለዓት የወቀ	
15	ያጠፋሉ?	(( <i>T</i> )A( <i>L</i> )		
P4	ሥራዎ በትንሹ ትንፋሽ ወይም የልብ ምት እንደ ድንነተኛ የእግር ጉተ	2. አያካትትም		
	ሸክሞችን] ቢያንስ ለ 10 ደቂቃዎች ያለማቋረጥ የሚጨምር መጠነኛ ኃይለኛ		3. ያካትታል	
	እንቅስቃሴን ያካትታል?			
P5	በተለመደው ሳምንት ውስጥ ሥራዎ መጠነኛ የአካል ብዋት እንቅስቃሴዎችን በስንት		0	
Dí	ቀናት ያካሂዳሉ?		<i>ቀ</i> ን	
P6	በተለመዳሡ ዋን በሥራ ሳይ መጠንና የለባል ብይተ አንዋቢቃቤዎተን ለማድረግ ምን ያህል ጊዜ ያጠፋሉ?		በሣተ <u>አዊ</u> ቃ	
P7	በተከታታይ ቢያንስ ለ 10 ደቂቃዎች በብስክሌት ይጓዛሉ ወይም ይጠቀ	2. አያካትትም		
		3. ያካትታል		
P8	በተለመደው ሳምንት ውስጥ ለምን ያክል ቀን በተከታታይ ቢያንስ ለ 10	ደቂቃዎች ስንት	<u></u> ۱	
Dô	ቀናት በእግር ወይም በብስክሌት ይጓዛሉ?	<i>Φ'</i>		
P9	በተለመደው ቀን ለጉዞ በእግር ወይም በብስክሌት ምን ያህል ጊዜ ያጠፋሪ	<u>ት?</u> መቀ እ ! = ፡ ነሻ	ሰዓተ ደቂቃ	
P10	እንደ [ሩጫ ወይም እግር ኳስ] ያለማቋረዋ ቢያንስ ለ 10 ደቂቃዎች ያለማቋረዋ ከፍተኛ		2. ለያባተተም 2. ውካ ከተ	
	፲ ነገገጥበ ወይንግ እናጠ ንግግ የማጨምር ፕግር ያጠበግሮ ሥይበት በፖ ር ብ ቀት እንቅስ ቀሰ ዎችን ጠይም የመዝናኛ (መዝናኛ) እንቅስ ቀሰ ዎችን የ(	<i>5. ዩ</i> ባተታል		

P11	ነተለመደው ሳምንት ውስጥ ኃይለኛ ስፖርቶችን ፤ የአካል ብቃት እንቅስቃሴዎችን		7		N		
	ወይም የመዝናኛ (መዝናኛ) እንቅስቃሴዎችን በስንት ቀናት ያካሂዳሉ?		ቀን				
P12	በተለመደው ቀን ጠንከር ያሉ ስፖርቶችን ፣ የአካል ብቃት እንቅስቃ	ተለመደው ቀን ጠንከር ያሉ ስፖርቶችን ፣ የአካል ብቃት እንቅስቃሴዎችን ወይም		ነትደ	<del>ጀ</del> ቃ		
	መዝናኛ እንቅስቃሴዎችን ለማድረግ ምን ያህል ጊዜ ያጠፋሉ?						
P13	ለማቋረጥ ለ 10 ደቂቃዎች በትንሹ ትንፋሽ ወይም የልብ ምት እንዲጨምር		2.	2. አያካትትም			
	የሚያደርጉ፣ [ብስክሌት መንዳት ፣ መዋኘት እና ቮሊቦል] መጠነኛ ኃይ	ሚያደርጉ፣ [ብስክሌት መንዳት ፣ መዋኘት እና ቮሊቦል] መጠነኛ ኃይለኛ ስፖርቶችን		3. ያካትታል			
	፤ የአካል ብቃት እንቅስቃሴዎችን ወይም መዝናኛ እንቅስቃሴዎችን ያደር.	<i>ጋ</i> ሉ?					
P14	በተለመደው ሳምንት ውስጥ መጠነኛ ኃይለኛ ስፖርቶችን ፤	የአካል ብቃት	•		<u> </u>		
	<u>እንቅስቃሴዎችን ወይም የመዝናኛ (መዝናኛ) እንቅስቃሴዎችን በስንት ቀ</u>	<u>ናት ያካሂዳሉ?</u>	ቀን	ቀን			
P15	ነተለመደው ቀን መጠነኛ ስፖርቶችን ፣ የአካል ብቃት እንቅስቃሴዎችን ወይም		՝ሰ٬	ሰዓትደቂቃ			
	የመዝናና (መዝናና) እንቀስቃሴዎትን ለማድረግ ምን ያህል ጊዜ ያጠፋሉ	?					
የመቀመ	ም ባህሪ						
S1	በተለመደው ቀን ለመቀመጥ ምን ያህል ጊዜ ያጠፋሉ?	ሰዓት	_ደቂቃ				
S2	በተለመደው ቀን ውስጥ ስንት ሰዓት ይተኛሉ፡፡	ትለ <i>መ</i> ደው ቀን ውስጥ ስንት ሰዓት ይተኛሉ፡፡ሰዓት					
<b>S</b> 3	በዙ ጊዜ እንቅልፍ <i>መ</i> ቅረጥ አለዎት? 1. የለም						
		2. አንዳንድ 1	"Њ				
		3.በአብዛኛው					
S4	በአንድ ወር ውስጥ ለምን ያህል ቀናት የጣታ ስራ ተረና እንዳለብዎ		በቀን				
	(ለሆስፒታል ሥራተኖተ ብቻ)						
የጭ	ንቀተ እና የጭንቀተ ልኬተ - 21 መለኪያ (DASS -21)						
- b ol							
7. AU	ነዎ <i>ን እያንጓንዱን መ</i> ባለጫ ያንተዘዙ እና ባለፈው ባምንተ ውበጥ መ	ግለጫው ለእር	ጠም <u>ም</u> ን ያ	ህል አንደ	ተተገበረ		
የሚ	ያመለክተ ቁጥር ሀ ፣ 1 ፣ 2 ወይም 3 ላይ ክበብ ያድርጉ።						
የደረጃ አሰጣጡ መጠን እንደሚከተለው ነው							
0 በጭራሽ በእኔ ላይ አልተተንበረም							
1 በተወሰነ ደረጃ ወይም በተወሰነ ጊዜ ለእኔ ተፈጻሚ ሆነ							
2 ለ	ኣኔ በተወሰነ ደረጃ ወይም በጥሩ ጊዜ ላይ ተፈጻሚ ሆነኝ						
3 ለ	ኣኔ ብዙ ወይም ብዙ ጊዜ አልተተገበረም						
1 (s)	እራሴን ለማደስታ እቸንራለው፡፡	0	1	2	3		
2 (a)	የአፈዎን ደረቅነት ያውቁ ነበር		1	2	3		
3 (d)	በጭራሽ ምንም አዎንታዊ ስሜት አይሰማኝም		1	2	3		
4(a)	የአተነፋፈስ ችግር አጋጥሞኛል (ለምዕሌ ፤ ከመጠን በለይ ፈጣን መተንፈስ ፤		1	2	3		
. (u)	አካላዊ ድካም በሌለበት መተንፈስ)		-	-	C		
5 (d)	ነገሮችን ለማድረግ ተነሳሽነቱን መሥራት ከባድ ይሆንብዎታል፡፡		1	2	3		
6 (s)	በሁኔታዎች ላይ ከመጠን በላይ አጸፋዊ ምላሽ እሰጣለው፡፡		1	2	3		
7 (a)	እንደ መንቀጥቀጥ አጋጥሞዎት ያውቅ ነበር (ለምሳሌ በእጆች )		1	2	3		
<b>8</b> (s)	ብዙ ጊዜ የመቆጣት ባህሪ የጠቀማሉ፡፡		1	2	3		
9 (a)	እራሴን ሞኝ የማደርግበት ሁኔታ አሳስቦኝ ያሳስበኛል፡፡		1	2	3		
10 (d)	ምንም የምጠብቀው ነገር እንደሌለኝ ይሰማኛል፡፡	0	1	2	3		
<b>11 (s)</b>	እኔ እራሴ ተበሳጭቼ አንኘዋለው፡፡		1	2	3		
12 (s)	እራሴን ዘና ለማድረግ እቸገራለው።		1	2	3		
13 (d)	የተስፋ መቁረጥ ስሜት ይሰማኛል፡፡		1	2	3		
14 (s)	እኔ የምሥራውን እንዳላደርግ የከለከለኝን ማንኛውንም ነገር አልታገስም		1	2	3		
15 (a)	ለድንጋጤ ቅርብ እንደሆንኩ ይስማኛል፡፡		1	2	3		

16 (d)	ስለማንኛውም ነገር ቅን መሆን አልቻልኩም ወይም አይሰማኝም፡፡		0	1	2		3
17 (d)	እንደ ሰው ብዙም ዋጋ እንደሌለኝ አይሰማኝም፡፡		0	1	2		3
<b>18</b> (s)	ዝም ብሎ እኔ የተነካው ይመስለኛል ፡፡		0	1	2		3
<b>19 (a)</b>	ምንም ነገር በሌለበት ፍራት ፍራት ይሰማኛል (ለምሳሌ የልብ ምት መል	ጌመር ፤	0	1	2		3
	የልብ ምት ማጣት)						
20 (a)	ያለ ምንም ምክንያት ፍርሃት ይሰማኝ		0	1	2		3
21 (d)	ሕይወት ትርጉም የለሽ እንደሆነ ይሰማኝ		0	1	2		3
የቤት ሁ	በት ፣ ቤተሰቡ ከሚከተሉት ንብረቶች ውስጥ አንዳቸውም አሉት?						
H1	የሲዲ ጣጫወቻ			0 9	ለም	1.	አለ
H2	የሚሰራ ጠፍጣፋ ማያ ገጽ ቴሌቪዥን	የሚሰራ ጠፍጣፋ ጣያ ነጽ ቴሌቪዥን			ለም	1.	አለ
H3	የ <i>ጋ</i> ዝ ምድጃ/ሲሊንደር			0 9	ለም	1.	አለ
H4	ማቀዝቀዣ ፍሪጅ			0 9	ለም	1.	አለ
H5	የኤሌክተሪክ ምድጃ			0 9	ለም	1.	አለ
H6	ባይክል			0 9	ለም	1.	አለ
H7	ምተር ሳይክል			0 9	ለም	1.	አለ
H8	26			0 9	ለም	1.	አለ
H9	ሰፍ			0 9	ለም	1.	አለ
H10	ፍራሽ			0 9	ለም	1.	አለ
H11	መኪና			0 9	ለም	1.	አለ
H12	ባጃጅ			0 9	ለም	1.	አለ
H13	ታክሲ			0 9	ለም	1.	አለ
H14	የባል ቤት			0 9	ለም	1.	አለ
H15	ታብሌት ኮምፒተር			0 9	ለም	1.	አለ
H16	ስማርት ስልክ			0 9	ለም	1.	አለ
H17	<i>ዲጂታ</i> ል ካሜራ			0 9	ለም	1.	አለ
H18	የልብስ ማጠቢያ ማሽን			0 9	ለም	1.	አለ
H19	ሳፕቶፕ			0 9	ለም	1.	አለ
H20	ካውያ			0 9	ለም	1.	አለ
በርጭ መጠን መለኪያ							
	ግብ መጠን 1ሴ.ሜ						
	<i>መ</i> ጠን 2			ሴ.ጣ	ļ		

#### **13. Declaration**

This is to certify that the proposal title entitled **central obesity and its associated factors among adults aged 18-64 years in Debre Tabor Town, northeast, Ethiopia, 2021,** submitted in partial fulfilment of the requirements for the degree of /Master of public health in human nutrition department of nutrition and dietetics, Bahir Dar University, is a record of original work carried out by me and has never been submitted to this or any other institution to get any other degree or certificates. The assistance and help I received during this investigation have been duly acknowledged.

Principal investigator			
Tigist Yismaw (BSc in PHO)			
Name	signature	Date	
Advisors			
Dr. Netsanet Fentahun(Double PhD,	Associate Prof)		
Name		Signature	Date
Ms. Hanna Demelash(MPH, Ass,t Pr	of)		
Name	Signature	Date	•