

2022-11-19

# Surgical Management Outcomes and Associated Factors of Epidural Hematoma in Tibebe Ghion Specialized Hospital and Felege Hiwot Comprehensive Specialized Hospital, Northwest Ethiopia

Aytenew, Amsalu

---

<http://ir.bdu.edu.et/handle/123456789/15034>

*Downloaded from DSpace Repository, DSpace Institution's institutional repository*



**BAHIR DAR UNIVERSITY**  
**COLLEGE OF MEDICINE AND HEALTH SCIENCES**  
**SCHOOL OF MEDICINE, DEPARTMENT OF Surgery**

**Surgical Management Outcomes and Associated Factors of Epidural Hematoma in  
Tibebe Ghion Specialized Hospital and Felege Hiwot Comprehensive Specialized  
Hospital, Northwest Ethiopia**

**By: - Dr Aytenew Amsalu (Md)**

**A THESIS RESEARCH SUBMITTED TO BAHIR DAR UNIVERSITY,  
COLLEGE OF MEDICINE AND HEALTH SCIENCES, SCHOOL OF  
MEDICINE AND DEPARTMENT OF SURGERY FOR PARTIAL  
FULFILLMENT OF SPECIALTY IN GENERAL SURGERY**

**NOVEMBER 2022**

**BAHIR DAR, ETHIOPIA**

**BAHIR DAR UNIVERSITY**

COLLEGE OF MEDICINE AND HEALTH SCIENCES  
SCHOOL OF MEDICINE, DEPARTMENT OF SURGERY

Investigator

Dr.Aytenew Amsalu (MD)

Phone: +251974497947

E-mail: [aytenew@gmail.com](mailto:aytenew@gmail.com)

Name and address of advisors

1. Melkamu Bedimo (Asso't Pro)

Phone: +251913221769

E-mail: [melkamubb@gmail.com](mailto:melkamubb@gmail.com)

2. Dr. Fisseha Gebeyehu (Ass't Pro, MD)

Phone: - +2519 512342

TITLE: SURGICAL MANAGEMENT OUTCOMES AND  
ASSOCIATED FACTORS OF EPIDURAL HEMATOMA AT TIBEBE  
GHION SPECIALIZED HOSPITAL AND FELEGE HIWOT  
COMPREHENSIVE SPECIALIZED HOSPITAL, NORTHWEST ETHIOPIA

NOVEMBER 2022

BAHIR DAR, ETHIOPIA

**Declaration**

This is to certify that the thesis entitled “Surgical Management outcomes and associated factors of epidural hematoma at Tibebe Ghion Specialized Hospital and Felege Hiwot Comprehensive

Specialized Hospital, Northwest Ethiopia, 2022” submitted in partial fulfillment of the requirements for the specialty in General surgery. The record of original work carried out by me and has never been submitted to Bahir Dar University or any other institution to get any other degree or certificates. The assistance and help I received during the course of this investigation have been duly acknowledged.

Dr Aytnew Amsalu (MD)

Name of the candidate

\_\_\_\_\_



Signature

\_\_\_\_\_

Date

### Approval of the advisors

I hereby certify that I have supervised, read, and evaluated this thesis titled "Surgical Management outcomes and associated factors of epidural hematoma at Tibebe Ghion Specialized Hospital and Felege Hiwot Comprehensive Specialized Hospital, Northwest Ethiopia, 2022" by Dr Aytenew Amsalu (MD) prepared under my guidance. I recommend the thesis be submitted for oral defense.

- |  |   |                   |
|--|---|-------------------|
| 1. <u>Melkamu Bedimo (MPH, Asso't pro)</u> | <u></u> | <u>9/3/25 E.C</u> |
| Advisor's name                             | Signature   | Date              |
| 2. <u>Dr. Fisseha Gebeyehu (Ass't Pro)</u> | <u></u> | <u>9/3/25 E.C</u> |
| Advisor's name                             | Signature   | Date              |



## **Acknowledgment**

First, I would like to acknowledge Bahir Dar University College of Medicine and Health Sciences, School of Medicine, department of Surgery for giving this opportunity to do.

Next, I would like to express my heart-felt gratitude to my advisors Melkamu Bedimo (MPH, Asso't pro) and Dr. Fisseha Gebeyehu (Ass't Pro), for their invaluable guidance and advice in preparation of this thesis research.

Furthermore, my gratitude goes to Tibebe Ghion Specialized Hospital and Felege Hiwot Comprehensive Specialized Hospital and data collectors for their unlimited support and giving the required information.

## Table of Contents

|  |                                     |
|--|-------------------------------------|
| Declaration .....                            | i                                   |
| Approval of the advisors .....               | <b>Error! Bookmark not defined.</b> |
| Acknowledgment .....                         | iv                                  |
| List of Table .....                          | vii                                 |
| List of figures .....                        | viii                                |
| List of annex .....                          | ix                                  |
| Abbreviations and acronyms.....              | xi                                  |
| Abstract .....                               | xii                                 |
| 1. INTRODUCTION .....                        | 1                                   |
| 1.1 Background .....                         | 1                                   |
| 1.2 Statement of problem .....               | 3                                   |
| 1.3 Significance of the study .....          | 5                                   |
| 2. LITERATURE REVIEW .....                   | 6                                   |
| 2.1 Outcomes of EDH.....                     | 6                                   |
| 2.2 Associated factors of EDH.....           | 7                                   |
| 2.2.1 Sociodemographic characteristics ..... | 7                                   |
| 2.2.2 Clinical related characteristics.....  | 7                                   |
| 2.2.3 Trauma related characteristics .....   | 8                                   |
| 2.3 Conceptual framework .....               | 10                                  |
| 3. Objectives .....                          | 11                                  |
| 3.1 General objective.....                   | 11                                  |
| 3.2 Specific Objective .....                 | 11                                  |
| 4. METHODS AND MATERIAL .....                | 12                                  |
| 4.1 Study area and period.....               | 12                                  |
| 4.2 Study design .....                       | 12                                  |
| 4.3 Population.....                          | 12                                  |
| 4.3.1 Source population .....                | 12                                  |
| 4.3.2 Study population.....                  | 12                                  |
| 4.4 Inclusion and Exclusion Criteria .....   | 12                                  |
| 4.4.1 Inclusion Criteria .....               | 12                                  |

|   |    |
|---|----|
| 4.4.2 Exclusion criteria.....                             | 12 |
| 4.5 Variables of the Study.....                           | 13 |
| 4.5.1 Dependent variables.....                            | 13 |
| 4.5.2 Independent variables.....                          | 13 |
| 4.6 Operational definition and term definitions.....      | 13 |
| 4.7 Sample size determination and sampling procedure..... | 13 |
| 4.8 Sampling technique and procedure.....                 | 14 |
| 4.9 Data collection tools and methods.....                | 15 |
| 4.10 Data quality control.....                            | 16 |
| 4.11 Data Processing and Analysis.....                    | 16 |
| 4.12 Ethical Consideration.....                           | 16 |
| 4.13 Dissemination of the Result.....                     | 17 |
| 5. Result.....  | 18 |
| 5.1 Socio-demographic characteristics.....                | 18 |
| 5.2 Clinical related characteristics.....                 | 19 |
| 5.4 Factors affecting EDH.....                            | 24 |
| 6. Discussion.....  | 25 |
| 6.1 Limitations of the study.....                         | 26 |
| 7. Conclusion and Recommendation.....                     | 27 |
| 7.1 Conclusions.....                                      | 27 |
| 7.2 Recommendation.....                                   | 27 |
| 8. Reference.....   | 28 |
| Annex.....  | 30 |



## List of Table

|   |    |
|---|----|
| Table 1:- Socio-demographic characteristics of the management outcome of EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022(n=366) .....            | 18 |
| Table 2:- Clinical presentation related characteristics of the management outcome of EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022(n=366)..... | 20 |
| Table 3:- CT scan findings of EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022(n=366) ...   | 21 |
| Table 4:- Type of management and outcome of EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022(n=366).....  | 23 |
| Table 5:- factors affecting the outcome of EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022(n=366).....   | 24 |

## List of figures

|  |    |
|--|----|
| Figure 1: Conceptual framework of epidural hematoma .....                                  | 10 |
| Figure 2: Schematic presentation.....  | 15 |
| Figure 3: Clinical presentation of the EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022..... | 19 |
| Figure 4: Mechanism injury of the EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022.....      | 21 |
| Figure 5: Site of EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022 .....                     | 22 |
| Figure 6: Treatment outcomes of EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022.....        | 23 |

**List of annex**

**Annex I: Information sheet** ..... 30  
Annex III: English Version Checklist..... 31



## **Abbreviations and acronyms**

|       |   |
|-------|---|
| CNS   | Central Nervous System                          |
| CT    | Computed Tomography                             |
| DM    | Diabetic Mellitus                               |
| DSF   | Depressed skull fracture                        |
| EDH   | Epidural Hematoma                               |
| FHCSH | Felege Hiwot Comprehensive Specialized Hospital |
| GCS   | Glasgow Coma Scale                              |
| ICP   | Intracranial pressure                           |
| IRB   | Institutional Review Board                      |
| RTA   | Road Traffic Accident                           |
| SOM   | School of medicine                              |
| SPSS  | Statistical Package for Social Sciences         |
| TBI   | Traumatic Brain Injury                          |
| TGTH  | Tibebe Ghion Teaching Hospital                  |
| WHO   | World Health Organization                       |
| MRI   | Magnetic Resonance Imaging                      |

## **Abstract**

**Background:** Epidural hematoma (EDH) is a major traumatic brain injury and a potentially life-threatening condition. Epidural hematoma is one of the most common causes of mortality and morbidity among head injury, but little is known about the problem in Northwest Ethiopia. Intracranial hematoma is the collection of blood in the various intracranial spaces.

**Objective:** To assess the surgical management outcome and associated factors of epidural hematoma in Felege Hiwot Comprehensive Specialized Hospital (FHCSH) and Tibebe Ghion Teaching Hospital (TGTH), Bahir Dar City, Northwest Ethiopia, 2022.

**Methods:** Institution-based retrospective cross-sectional study among 366 epidural hematoma patients charts from January 2020 January 20, 2022 at TGTH and FHCSH, Northwest Ethiopia. Systematic random sampling technique was used to select patient charts. Data were collected by reviewing patient charts using pre-tested questionnaires. Data were entered into the Epi data version 3.1 and exported to SPSS version 23 for further analyses. A binary logistic regression model was fitted to identify the associated factors. Variables with P-value < 0.25 in the bi-variable analysis were a candidate for multivariable analysis and P-value < 0.05 in the multivariate analysis were used to declare as statistically significant. The odds ratio (OR) with a 95% confidence interval (CI) was used to measure the strength of association.

**Results:** Over all, death among EDH treatment was 20(5.5%, 95% CI: 3.3-7.7%) outcome was death. The most common presentation of EDH were headache 343(93.7%), LOS 330(90.2%), vomiting 298(81.4%), and 107(10.9%) increased ICP. Of patients with increased ICP 64(59.8%) had decreased mentation followed by 33(33.7%) vomiting and 91(24.9%) had hypertension. The increased ICP (AOR=5.72, 95% CI: 1.55-21.07), >25mm thickness of hematoma (AOR=5.29, 95% CI: 1.43-19.61) and hypotension (AOR=4.30, 95% CI: 1.41-13.08) was significantly associated with the outcomes of EDH treatment.

**Conclusion:** The outcome of EDH treatment was comparable with other studies. The increased ICP, >25mm thickness of hematoma and hypotension was significantly associated with the outcomes of EDH treatment. Health care professionals should closely monitor blood pressure and the size hematoma >25mm.

Key words: Epidural hematoma, traumatic brain injury, outcome, trauma, craniotomy

# 1. INTRODUCTION

## 1.1 Background

An epidural hematoma (EDH) is a collection of blood that forms between the skull and the dura mater, the outermost protective membrane covering the brain. It is bleeding between the inside of the skull and the outer covering of the brain (called the dura) and includes collection of blood in epidural, subdural, parenchymal and intraventricular areas. Other names for EDH are extradural hematoma, epidural hemorrhage or intracranial hematoma. More broadly, it is also a type of traumatic brain injury (TBI). It is a life-threatening complication of severe traumatic brain injury (TBI)(1). It may be classified to into three groups: traumatic epidural hematoma which is the main cause of epidural hematoma, spontaneous spinal epidural hematomas and postoperative epidural spinal hematoma(1, 2).

Cranio-cerebral lesions can be classified into primary and secondary. Primary lesions occur as a direct result of a trauma to the head and include scalp injuries, skull fractures, extra-axial hemorrhage and intra-axial lesions. Secondary brain injury occurs as a complication of primary lesions and includes ischemic and hypoxic damage, cerebral edema and brain herniation. Extra-axial hemorrhage includes EDH, subdural hematoma (SDH), and subarachnoid (SAH) and intraventricular (IVH) hemorrhage. Primary intra-axial injuries include diffuse axonal injuries (DAIs), cortical contusions, intra-parenchymal hematomas (IPHs) and vascular lesions(3).

It is a potentially life-threatening complication of a head injury and usually caused by a head injury. The most common cause of EDH is trauma, microscopic tears in the blood vessels under the skin, bleeding disorders, and thinner skin due to aging(4). A skull fracture occurs in 75% of the cases. Physical abuse, vehicular accident, or collision in contact sports can also cause head injury and lead to an epidural hematoma. An EDH is diagnosed using computed tomography (CT) or magnetic resonance imaging (MRI) of head or spine. In the head, a hematoma appears as a dense mass that pushes the brain away from the skull.

The clinical symptoms can also occur in mild to moderate TBI and EDH can develop rapidly after an injury or slowly over the course of several hours. The time it takes for symptoms to develop depends on the severity of the injury and how quickly blood is filling into the lining

between the brain and skull. Some of the most common signs and symptoms of an EDH include; vomiting, confusion, seizures, loss of vision on one side, dizziness, changes in breathing, nausea, severe headache, weakness in one half of the body, and drowsiness or loss of alertness. The possible complications of EDH are herniation of the brain and permanent coma, normal pressure hydrocephalus, and paralysis or loss of sensation (which began at the time of the injury)(5).

Management of EDH is both conservative and surgical. Conventionally, urgent evacuation is the definitive mode of management(6). With the routine use of CT non-operative management is being used more often in selected patients(7-9) Evacuation of the hematoma, coagulation of bleeding sites, and inspection of the dura follows the craniotomy and burr hole(10, 11). The decision to perform a surgery in a patient with a traumatic EDH is dependent on several factors (neurologic status, size of hematoma, age of patients, CT findings). Indications for surgery: An EDH greater than 30 cm<sup>3</sup> should be surgically evacuated regardless of the patient's Glasgow Coma Scale (GCS) score. To date, the outcomes of AEDH vary from center to center, depending on the resources and quality of care in different regions(12). The management outcome of EDH may be measured using; morbidity, mortality, functional status, mental wellbeing, and other aspect of health-related quality of life (13-15).



## 1.2 Statement of problem

Annually 69 million people suffer from TBI, of whom 5.48 million experience severe TBI(16). Cranial EDHs result from TBI and represent a potentially life-threatening neurosurgical emergency(17). It can be associated with significant morbidity and mortality if untreated; however, patients with isolated EDH who are fortunate enough to undergo safe and timely surgical evacuation often sustain favorable outcomes(18).

A systematic review and meta-analysis study shows that, an overall EDH incidence of 8.2 % and an operative rate of 55.5 %, 3.1 million people worldwide require surgery for traumatic EDH every year, most of whom are in prime working age. Given the favorable prognosis with treatment, traumatic EDH is a strong investment for neurosurgical capacity building(19). An EDH can be a life-threatening condition and results in death in up to 15% of cases. It usually requires immediate treatment or can cause brain damage or possibly death if left untreated. Injuries continue to be an important cause of morbidity and mortality in the developed and developing world(20).

Study performed in the United States has estimated that EDH occurs in approximately 3 % of head injury cases(20). The study in Australia, indicate that the Injury Severity Score adjusted in-hospital mortality rates for young patients with EDH were 4.8%(21).

An EDH occurs in 2% of all head injuries and up to 15% of all fatal head traumas. Males are more often affected than are females. Furthermore, the incidence is higher among adolescents and young adults. The mean age of affected patients is 20 to 30 years, and it is rare after 50 to 60 years of age. As an individual's age advances, the dura mater becomes more adherent to the overlying bone. This decreases the chance that a hematoma can develop in the space between the cranium and dura(22)

Trauma is the leading cause of death in people younger than 45yrs worldwide. <sup>(1)</sup> There is evidence that prompt neurosurgical management of TBI can significantly improve outcome, especially if decompression is performed within 48 h of injury(23). An EDH occurs in 2% of all head injuries and up to 15% of all fatal head traumas(22). EDHs occur in 0.2–12% of acute head-injured patients, and the overall mortality is 5%(24). EDHs generally occur at the site of impact, and a fracture is associated in 90% of cases(25).

It is a life-threatening condition, which may require immediate intervention and can be associated with significant morbidity and mortality if left untreated. Rapid diagnosis and evacuation are important for a good outcome (13, 26).

According to previous studies done in Ethiopia, the prevalence of head injury in patients presenting to the surgical emergency department ranged from 24.9% to 49.4% (27-29). More than half of the patients sustained a mild head injury 53.3% and 63.8% an open head injury(29).

A study in Hawassa shows that, the incidence of the mortality rate in TBI was 2.26 per 100-person day observation. The independent predictors of time to death were age above 65 years severe TBI, moderate TBI, hypotension, hypoxia, hyperthermia and hypoglycemic positively associated with mortality, while underwent neurosurgery was negatively associated with mortality(30). In Addis Ababa the prevalence of acute EDH was 15.9% which is the second leading cause of head trauma injury admission(31).

Generally, in our centers (FHCSH and TGTH) patients with AEDH are one of the commonly encountered patients but our health care area doesn't have well-structured document, and full information on management outcome that creates difficulty of showing health care improvement. Doing research on management outcome provides realistic information about benefit and risk of specific procedure and guides institution and addresses their problem. In Ethiopia, population and patient visits to hospitals that provide or doing surgical procedure are increasing in number.

### **1.3 Significance of the study**

This study can also help surgical staff in the setting an opportunity to assess their performance and a chance to improve the outcomes of epidural hematoma. The hospital admin can also be made to give a special attention on surgical service provision to improve the management outcome.

This study is important to provide crucial information about the predisposing factors, clinical characteristics and outcome of epidural hematoma. The finding of this study will be identifying the factors of epidural hematoma. This should help the physician and the patient to choose better treatment options for reduction of morbidity, mortality and cost of health care.

The findings of this study will be used as inputs for program implementers at national as well as regional levels to improve treatment outcomes of epidural hematoma. This study can be used to guide a clinician for choosing the appropriate managements of epidural hematoma. Therefore, this study used as an input for other researchers to do further analysis.

## **2. LITERATURE REVIEW**

### **2.1 Outcomes of EDH**

A study in Pakistan shows that 62.4% EDH patient's outcome was good and predominantly these patients were in GCS 14 – 15 on presentation. Death occurred in 7.90% patients and 0.817% patients remained in vegetative state. These patients had very low GCS and coning or associated parenchymal or systemic injuries were already present before their operations(12). A study in Nigeria shows that the mortality rate of patients presented with EDH was 16.9%.(32)

A study in Mulago National Referral Hospital, Uganda shows that overall proportion of favorable outcome in patients with extra axial hematoma was 71.7%, with 42.3% and 81.7% for ASDH and AEDH, respectively. Factors associated with a favorable outcome were admission systolic BP > 90 mmHg, oxygen saturation > 90% and diagnosis AEDH. Moderate TBI and severe TBI were significantly associated with unfavorable outcomes(33).

A study in India shows that about 70% of patients recover well, 20% had moderate disability, 3.33% had severe disability, 3.33% were in persistent vegetative state, and 1 3.33% patient were dead(34). Although conservative management is successful management of epidural hematoma, surgical evacuation constitutes definitive treatment(9).

A study in Australia shows that, 21.5 % of severe TBI had EDH; of these, 30.8 % died in the hospital, 13.2 % survived with unfavorable outcome, 51.6 % with favorable outcome. Mortality rates predicted by the Rotterdam score showed good correlation with observed mortality rates(35). Study show that patients with EDH who have a high GCS score 13–15, volume <40 mm, and less than 6 mm of midline shift should be considered for conservative management(36).

A study in Hawassa shows that, 72.2% TBI cases were managed conservatively, while 27.8% underwent various surgical interventions within the first one week of presentation. Most of the operated cases were aged 5 to 10 years (55%). The common surgical indications were the evacuation of epidural and subdural hematoma (4.5%); wound debridement for compound skull fracture (21.3%); and depressed skull fracture elevation (74.2%). Burr hole and evacuation was done for four mild TBI cases(37).

## **2.2 Associated factors of EDH**

### **2.2.1 Sociodemographic characteristics**

A study in Pakistan shows that, majority of patients with EDH was younger than 60 years; and the commonest age group was 21 to 30 years(27.79%) (12). A study in India shows the largest number of patients having head injury belong to age group of 21-30 years (50%), followed by 41-50 years age group (16.66%), less common in age group more than 60 years (3.3%). Male to female ratio was 4:1(34). Males are more often affected than females, and the incidence is higher among adolescents and young adults. The mean age of affected patients is 20 to 30 years, and it is rare after 50 to 60 years of age. As an individual's age advances, the dura mater becomes more adherent to the overlying bone. This decreases the chance that a hematoma can develop in the space between the cranium and dura(22).

A study in Australia shows that, age was the main factors influencing outcomes after severe TBI associated with EDH(35).

A study in Hawassa shows that, about 79.8% of falls and 58.3% of RTAs occurred in boys(37).

A study in Gedeo Zone, Southern Ethiopia; the majority of head injury patients were found in the age categories of 15-29 years (36.8%), and under 15 years (29.2%). Majority of patients were male dominance of head injury patients (71.7%)(27).

### **2.2.2 Clinical related characteristics**

A study in Pakistan shows that, the commonest presentation of EDH was 73.11% headache, 63.44% had vomiting, 32.7% loss of consciousness and 26.97% focal neurological deficit as presenting complaints. The site of hematoma was temporal-parietal(30.79%), frontal(26.97%), parietal(12.8%), posterior fossa (2.99%), and occipital (2.99%) (12).

A study in India shows the mode of injury in majority of patients was due to road traffic accidents (60%), self-fall (30%), and assault (10%). Patients presented with lucid interval (36.66%), anisocoria (23.33%), Battle's sign (13.33%), hemotympanum (10%), black eye (10%), and hypoxia or hypotension (6.66%). Majority of mode of EDH management 63.33% were conservatively (34).

A study in Australia shows that, severity of TBI, and neurological status were the main factors influencing outcomes after severe TBI associated with EDH(35).

Study show that a low GCS before surgery, or on arrival, abnormal pupil examination, the time between neurological symptoms and surgery, and elevated ICP in the post-operative period were associated with poor prognosis of EDH. The typical presentation is an initial loss of consciousness following trauma, a complete transient recovery (“often termed as a lucid interval”), culminating in a rapid progression of neurological deterioration. This occurs in 14% to 21% of patients with an EDH. Therefore, the presentations range from a temporary loss of consciousness to a coma. The presentation of symptoms depends on how quickly the EDH is developing within the cranial vault. A patient with a small EDH may be asymptomatic, but this is rare. Also, an EDH may also develop in a delayed fashion. Enlarging hematoma leads to eventual elevation of intracranial pressure which may be detected in a clinical setting by observing ipsilateral pupil dilation, the presence of elevated blood pressure, slowed heart rate, and irregular breathing(1).

A study in Hawassa shows that TBI contributed to 7.4% of pediatric emergency room visits. Pedestrian RTA and falls, early presentation (<24 hours of injury), and mild form of TBI among boys were the most common documented patterns. ICP, hyperglycemia, severe TBI, and presence of contusion, DAI, or intracranial bleeding on head CT predicted poor outcome. Majority (34%) of the TBI cases stayed from 4 to 7 days, while 78 (24.6%) were discharged within 24 hours of arrival. The average length of hospitalization days among those who died was 4.5 days (median 2 days), and 50% of deaths occurred within the first 3 days of admission. Factors affecting the outcomes were presence of increased ICP at admission, severe TBI, presence of hyperglycemia and contusion, DAI, or intracranial bleeding on head CT scan found to be significantly associated with death or neurologic disability among pediatric TBI(37).

### **2.2.3 Trauma related characteristics**

A study in Pakistan shows that location of epidural hematoma is significant because the prognosis relies on location of EDH. Supra tentorial epidural hematomas are more common than posterior fossa epidural hematomas. The most common location for EDH was temporo-parietal region 30.79% followed by frontal region 26.97%(12).

A study in India shows the most common site of EDH was temporal (20%), and temporo-parietal (20%) regions, parietal (16.66%) and frontal (13.33%) regions. About 56.6% had EDH on right side, 36.66% on left side and 6.66% had Bilateral EDH. CT brain findings of EDH shows; 46.66% patients clot volume was more than 30ml, and 53.33% of patients was found in less than 30ml. Midline shift was noted in 40% of patients, absent in 60% patients, and brain herniation was absent in 93.33% patients (34).

A study in Hawassa shows that, the main cause of TBI was RT (45.4%), falls (32.8%); fighting/violence (12.6%); and animal bite or kick injury (8.8%)(37).

## 2.3 Conceptual framework




Figure 1: Conceptual framework of epidural hematoma



### **3. Objectives**

#### **3.1 General objective**

- ✓ To assess the Surgical management outcome and associated factors of epidural hematoma in FHCSH and TGTH, Northwest Ethiopia 2022

#### **3.2 Specific Objective**

- To identify the surgical management outcomes of EDH in TGTH and FHCSH.
- To identify the associated factors of EDH in TGTH and FHCSH

## **4. METHODS AND MATERIAL**

### **4.1 Study area and period**

This study was carried out in Tibebe Ghion Teaching Hospital (TGTH) and Felege Hiwot Comprehensive Specialized Hospital (FHCSH), from August 1-30/2022. TGTH and FHCSH found in Bahir Dar city, is the capital city of Amhara regional state, found in northwest Ethiopia, 565 Km far from Addis Ababa; the capital city of Ethiopia.

Bahir Dar city have two public Comprehensive Specialized hospitals. This study was conducted in TGTH and FHCSH from October 15-November 13/2022 G.C. since it is possible to obtain a sufficient number of surgical patients coming from different parts of Northern Ethiopia; Department of neurosurgery is one of the main departments in TGTH and FHCSH, which gives full-fledged clinical service.

### **4.2 Study design**

Institution-based retrospective cross-sectional study design was conducted

### **4.3 Population**

#### **4.3.1 Source population**

Surgical patients with epidural hematoma from January 2020 to January 2022.

#### **4.3.2 Study population**

All patients diagnosed and surgically treated for Epidural hematoma.

### **4.4 Inclusion and Exclusion Criteria**

#### **4.4.1 Inclusion Criteria**

All surgical patients with epidural hematoma treated surgically were included. All patients who had a brain injury requiring admission and patients fit for operative modes in the study period were included.

#### **4.4.2 Exclusion criteria**

Those patients referred out with post-craniotomy complications were excluded. Patients with incomplete documentation and unknown discharge results were excluded.

## 4.5 Variables of the Study

### 4.5.1 Dependent variables

- ❖ Management (surgical) outcomes of EDH

### 4.5.2 Independent variables

- ❖ Socio-demographic characteristics including; sex, age, and residence
- ❖ **Clinical related characteristics including; mode of management, duration before operation, GCS during admission, vital sign at admission, investigation, comorbid illness, and anesthesia type**
- ❖ **Trauma related variables:** mechanism of injury, severity of injury, size of epidural hematoma, concomitant injury, urgency of surgery, and duration of presentation.

## 4.6 Operational definition and term definitions

**Outcome:** the final result of surgical patient after surgical management (at discharge)

**Emergent surgery:** surgery for a condition which is immediately life-threatening. Surgery must be performed within a few hours

**Urgent surgery:** surgery for a condition that is potentially life-threatening surgery usually must be completed within 24 hours

**Co morbidity:** the presence of one or more additional disease or disorder co-occurring with a primary disease or disorder

**Anesthesia type:** category that enables the painless performance of surgery.

**Complication:** is unfavorable evolution or consequence of a disease, a health condition or a therapy.

**Craniotomy:** it is a procedure of open up part of skull and removal of an epidural hematoma to reduce the pressure on the brain.

**Burr hole:** Drilling a small hole in the skull to relieve pressure and allow blood to drain outside the skull.

## 4.7 Sample size determination and sampling procedure

The sample size was determined using a single population proportion formula by considering a 95% confidence level, 5% marginal error and 39.0% the proportion of EDH in Addis Ababa(31).

Where;

n=desired sample size

$Z_{\alpha/2}$ = the value of standard score at 95% confidence level (1.96),

P= proportion of EDH (39.0%)

d= marginal error= 5% (0.05)

$$n_i = \frac{(Z_{\alpha/2})^2 PQ}{d^2} = \frac{(1.96)^2 (0.39 * (1 - 0.39))}{0.05^2} = \frac{0.91392}{0.0025} = 365.57 \approx \underline{\underline{366}}$$

Therefore, the final sample size is 366.

#### **4.8 Sampling technique and procedure**

In Bahir Dar city there are two public hospitals providing epidural hematoma management. Two hospitals were included based on adequate information it has been provided and its number of patients flow for the study area. A simple random sampling technique was used to select patients chart from the surgeon's registration log book. A patient's chart with incomplete documentation and unknown discharge result were replaced with the next charts.

|                |
|----------------|
| Bahir Dar city |
| 2745           |




Figure 2: Schematic presentation

#### 4.9 Data collection tools and methods

Data were collected by chart review retrospectively using structured pre-tested questioners and checklist from the patients' charts. The English language version questioners and checklist were used that adapted from different pieces of literature. The questionnaire and checklist contains; socio-demographic and clinical characteristics. The data were collected by two trained data collectors and supervised by the principal investigator.

#### **4.10 Data quality control**

Before the actual data collection, data collectors and supervisors were trained intensively on the contents of the questionnaire, checklist, data collection methods, and ethical concerns. Therefore, data collectors become familiar with the questionnaire.

Five percent of the samples were pre-tested in FHCSH and the questionnaire was modified and edited based on the findings. During data collection, both supervisors and the principal investigator were have checked the data for its completeness and missing information at each point. Furthermore, data were checked during entry and compilation before analysis.

#### **4.11 Data Processing and Analysis**

The collected data were coded, entered into EPI data version 3.1 software, and exported to statistical product and service solutions (SPSS) version 23 software packages for further analysis. Descriptive statistics were used to summarize the data in the form of frequency, mean, standard deviation (SD), and cross-tabulation. A binary logistic regression analysis was carried out to identify the association between the outcome variable and independent variables.

Variables with P-value  $<0.25$  in bi-variable logistic regression were used for multivariable logistic regression. A p-value  $<0.05$  were considered statistically significant. Adjusted odds ratio (AOR) with 95% confidence intervals (CI) was used.

#### **4.12 Ethical Consideration**

Ethical clearance was obtained from the Institutional Review Board (IRB) of the College of Medicine and health sciences, Bahir Dar University, Ethiopia. Then legal official clearance letters were written to the Ethiopian public health institute (EPHI). Finally, legal official clearance letters were written for TGTH and FHCSH.

Confidentiality was secured by avoiding writing the patient's name and the data were not give to the third person. The data extractions were conducted in a separate room. Moreover, confidentiality was secured during data collecting, analyzing, and reporting.

#### **4.13 Dissemination of the Result**

The plan of dissemination of the research result includes a presentation at Bahir Dar University College of Medicine and Health Sciences, Research Conferences. The report paper will also be disseminated to, Bahir Dar city health administration and regional health bureau. Publication in Scientific journal and online dissemination will also be considered.

## 5. Result

### 5.1 Socio-demographic characteristics

A total of 366 patient charts were included in the study. The mean age was  $29.95 \pm 11.42$  years (Mean  $\pm$  SD) with a range of 2-63 years. About 248(67.8%) of patients were rural residents, 327(89.3%) were male, 214(58.5%) of patients were 16-30 years and 219(59.8) of patients were farmers (Table1).

Table 1:- Socio-demographic characteristics of the management outcome of EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022(n=366)

| Variables  | Categories      | Frequency | Percentage |
|------------|-----------------|-----------|------------|
| Residence  | Rural           | 248       | 67.8       |
|            | Urban           | 118       | 32.2       |
| Sex        | Male            | 327       | 89.3       |
|            | Female          | 39        | 10.7       |
| Age        | $\leq 15$ years | 20        | 5.5        |
|            | 16-30years      | 214       | 58.5       |
|            | 31-45years      | 94        | 25.7       |
|            | $>45$ years     | 38        | 10.4       |
| Occupation | Farmer          | 219       | 59.8       |
|            | Merchant        | 57        | 15.6       |
|            | Driver          | 30        | 8.2        |
|            | Gov't employ    | 27        | 7.4        |
|            | Others          | 33        | 9.0        |



## 5.2 Clinical related characteristics

This study finding shows that majority of patients were presented with headache 343(93.7%), LOS 330(90.2%), vomiting 298(81.4%), and 107(10.9%) increased ICP. Of patients with increased ICP 64(59.8%) had decreased mentation followed by 33(33.7%) presented with vomiting and 91(24.9%) had hypertension. However, hypotension 40(10.9%), aspiration 30(8.2%) and seizure 25(6.8%) were less common clinical presentation of EDH (Figure 2 and Table 2).




Figure 3: Clinical presentation of the EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022

Ten (2.7%) of patients had chronic illness, of this six and four patients had hypertension and respiratory disease respectively. Only 6(1.6%) of patients had history of medication, and 45(12.3%) of patients had abnormal body moment. About 270 (73.8%) of patients place of occurrence trauma was outdoor. 153 (41.8%) of patients who had body weakness were hemiparetic. 298(81.4%) were right handed. Majority, 250(68.3%), of patients arrived with in

24hours. More than half 208(56.8%) patients had moderate TBI, 46(12.6%) of severe TBI and 6(1.6%) were critical head injury patients (Table 2).

Table 2:- Clinical presentation related characteristics of the management outcome of EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022(n=366)

| Variables                 | Categories                  | Frequency | Percentage |
|---------------------------|-----------------------------|-----------|------------|
| Chronic illness           | Yes                         | 10        | 2.7        |
|                           | No                          | 356       | 97.3       |
| Type of chronic illness   | hypertension                | 6         | 60.0       |
|                           | respiratory disease         | 4         | 40.0       |
| History of medication     | Yes                         | 6         | 1.6        |
|                           | No                          | 360       | 98.4       |
| Duration of hospital stay | ≤3days                      | 48        | 13.1       |
|                           | 4-7 days                    | 242       | 66.1       |
|                           | >7 days                     | 76        | 20.8       |
| Place of occurrence       | Occupant vehicle            | 36        | 9.8        |
|                           | Pedestrians                 | 18        | 4.9        |
|                           | Outdoor                     | 270       | 73.8       |
|                           | Home                        | 42        | 11.5       |
| Arrival time              | ≤12 hrs                     | 258       | 43.2       |
|                           | 23-24 hrs                   | 91        | 24.9       |
|                           | >24hrs                      | 117       | 32         |
| Abnormal body moment      | Yes                         | 45        | 12.3       |
|                           | No                          | 321       | 87.7       |
| Dominance                 | Right handed                | 298       | 81.4       |
|                           | Left handed                 | 68        | 18.6       |
| GCS                       | Mild                        | 161       | 44.0       |
|                           | Moderate                    | 134       | 36.6       |
|                           | Severe                      | 71        | 19.4       |
| Increased ICP             | Yes                         | 107       | 29.2       |
|                           | No                          | 259       | 70.8       |
| Symptoms of Increased ICP | Decreased mentation         | 64        | 59.8       |
|                           | Vomiting                    | 36        | 33.7       |
|                           | Hypertension                | 7         | 6.5        |
| Pupillary sign            | Midsized & reactive         | 185       | 50.5       |
|                           | Unilaterally dilated &fixed | 143       | 39.1       |
|                           | Bilaterally dilated &Fixed  | 38        | 10.4       |
| body weakness             | Hemiparesis                 | 153       | 41.8       |
|                           | Monoparesis                 | 27        | 7.4        |
|                           | Hemiplegia                  | 40        | 10.9       |
|                           | No                          | 146       | 39.9       |
| Severity of TB            | Mild                        | 106       | 29.0       |

|  |                 |     |      |
|--|-----------------|-----|------|
|  | moderate        | 208 | 56.8 |
|  | Severe/critical | 52  | 14.2 |

From the mechanism of injury fighting is the leading cause 72.7% and fall down injury was the second cause of EDH (Figure 4).




Figure 4: Mechanism injury of the EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022

### CT scan findings

The most commonly involved hemisphere was left 213(58.2) and 273(74.6%) of patients had more than 5cm midline shaft and the CT scan shows that 156(42.6%) of patients had linear skull fracture and 106(29.0%) patients had contusion (Table 3).

Table 3:- CT scan findings of EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022(n=366)

| Variables                 | Categories | Frequency | Percentage |
|---------------------------|------------|-----------|------------|
| Thickness of hematoma     | ≤25mm      | 224       | 61.2       |
|                           | >25mm      | 142       | 38.8       |
| Involvement of Hemisphere | Right      | 148       | 40.4       |
|                           | Left       | 213       | 58.2       |
|                           | Bilateral  | 5         | 1.4        |
| Midline shaft             | No         | 10        | 2.7        |
|                           | ≤5cm       | 83        | 22.7       |
|                           | >5cm       | 273       | 74.6       |
| Other CT scan findings    | None       | 44        | 12.0       |

|  |                       |     |      |
|--|-----------------------|-----|------|
|  | Linear skull fracture | 156 | 42.6 |
|  | DSF                   | 56  | 15.3 |
|  | Contusion             | 106 | 29.0 |
|  | others                | 4   | 1.1  |

The most common site of EDH was Parieto-temporal 196(53.6%), followed by frontal 89(24.3%), temporal 57(15.6%) and 24(6.5%) other sites (Figure 5).




Figure 5: Site of EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022

For majority of patients ,346(94.5%), craniotomy was done, 354(96.7%) were given general anesthesia and 64(17.5%) of patients were admitted in the ICU. The most common cause of death was primary brain injury 16(80.0) and respiratory failure accounted 4(20.0) (Table 4)

Table 4:- Type of management and outcome of EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022(n=366)

| Variables                  | Categories                  | Frequency | Percentage |
|----------------------------|-----------------------------|-----------|------------|
| Type of surgical procedure | Craniotomy                  | 346       | 94.5       |
|                            | Elevation & evacuation      | 9         | 2.5        |
|                            | Burr hole                   | 11        | 3.0        |
| Type of anesthesia used    | Local anesthesia            | 12        | 3.3        |
|                            | General anesthesia          | 354       | 96.7       |
| ICU admission              | Yes                         | 64        | 17.5       |
|                            | No                          | 302       | 82.5       |
| Cause of death             | Primary brain injury        | 16        | 80.0       |
|                            | Respiratory failure         | 4         | 20.0       |
| Neurologic outcome         | Good                        | 199       | 57.5       |
|                            | Some disability             | 128       | 37.0       |
|                            | Severe disability           | 15        | 4.3        |
|                            | persistent vegetative state | 4         | 1.2        |

About 20(5.5%, 95% CI: 3.3-7.7%) EDH treatment outcome was death while 346(94.5%, 95% CI: 92.3-96.7%) patients were discharged improved (Figure 6).




Figure 6: Treatment outcomes of EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022

## 5.4 Factors affecting EDH

On bi-variable analysis; residences, increased ICP, hypotension, duration of hospital stay, thickness of hematoma, severity of TBI, and time of arrival were associated with the outcomes of EDH treatment. On multivariable analysis; increased ICP, >25mm thickness of hematoma and hypotension was significantly associated with the outcomes of EDH treatment.

Hypotensive patients (systolic blood pressure less than 90mmHg) were 4.30 times more likely to die as compare to those with blood pressure greater than 90mmHg (AOR=4.30, 95% CI: 1.41-13.08)

Those with increased ICP EDH patients were 5.72 times more likely to die as compare to whose ICP was normal (AOR=5.72, 95% CI:1.55-21.07).

Those with >25mm thickness of hematoma patients were 5.29 times more likely to die as compare to whose thickness of hematoma ≤25mm (AOR=5.29, 95% CI: 1.43-19.61)(Table 6).

Table 5:- factors affecting the outcome of EDH at TGTH & FHCSH, Northwest, Ethiopia, 2022(n=366)

| Variables                    | Categories      | EDH                |                  | COR with 95% CI         | AOR with 95% CI         | P-value      |
|------------------------------|-----------------|--------------------|------------------|-------------------------|-------------------------|--------------|
|                              |                 | Improved           | death            |                         |                         |              |
| Residence                    | Rural           | 237(95.6%)         | 11(4.4%)         | 0.56(0.23-1.40)         | 0.48(0.16-1.45)         | 0.190        |
|                              | Urban           | 109(92.4%)         | 9(7.6%)          | 1                       | 1                       |              |
| <b>Increased ICP</b>         | <b>Yes</b>      | <b>92(86.0%)</b>   | <b>15(14.0%)</b> | <b>8.28(2.93-23.43)</b> | <b>5.72(1.55-21.07)</b> | <b>0.009</b> |
|                              | <b>No</b>       | <b>254(98.1%)</b>  | <b>5(1.9%)</b>   | <b>1</b>                | <b>1</b>                |              |
| <b>Thickness of hematoma</b> | <b>≤25mm</b>    | <b>220(98.20%)</b> | <b>4(1.8%)</b>   | <b>1</b>                | <b>1</b>                | <b>0.013</b> |
|                              | <b>&gt;25mm</b> | <b>126(88.7%)</b>  | <b>16(11.3%)</b> | <b>8.28(2.93-23.43)</b> | <b>5.29(1.43-19.61)</b> |              |
| Duration of hospital stay    | ≤3 days         | 43(89.6%)          | 5(10.4%)         | 1.36(0.39-4.72)         | 0.87(0.15-5.22)         | 0.883        |
|                              | 4-7 days        | 233(96.3)          | 9(3.7)           | 0.45(0.16-1.31)         | 0.92(0.23-3.65)         | 0.905        |
|                              | 7days           | 70(92.1%)          | 6(7.9%)          | 1                       | 1                       | 0.987        |
| Severity of TB               | Mild            | 101(95.3%)         | 5(4.7%)          | 1                       | 1                       | 0.081        |
|                              | moderate        | 203(97.6%)         | 5(2.4%)          | 0.50(0.14-1.76)         | 0.23(0.05-1.10)         | 0.065        |
|                              | Severe/Critica  | 42(80.8%)          | 10(19.2%)        | 4.81(1.55-14.92)        | 0.87(0.18-4.28)         | 0.863        |
| Arrival time                 | ≤12hrs          | 146(92.4%)         | 12(7.6%)         | 1                       | 1                       | 0.168        |
|                              | 13-24 hrs       | 85(93.4%)          | 6(6.6%)          | 0.86(0.31-2.37)         | 1.86(0.53-6.58)         | 0.336        |
|                              | >24 hrs         | 115(98.3%)         | 2(1.7%)          | 0.21(0.05-0.96)         | 1                       | 0.183        |
| <b>Hypotension</b>           | <b>Yes</b>      | <b>78(85.7%)</b>   | <b>13(14.2%)</b> | <b>6.38(2.46-16.55)</b> | <b>4.30(1.41-13.08)</b> | <b>0.010</b> |
|                              | <b>No</b>       | <b>268(97.5%)</b>  | <b>7(2.5%)</b>   | <b>1</b>                | <b>1</b>                |              |

*Hosmer and Lemeshow Test (P=0.070)*

## 6. Discussion

This study included 366 participants with a response rate of 100%. This study showed that 20(5.5%, 95% CI: 3.3-7.7%) of EDH patients were died. The study identified that increased ICP (AOR=5.72, 95% CI: 1.55-21.07), >25mm thickness of hematoma (AOR=5.29, 95% CI: 1.43-19.61) and hypotension (AOR=4.30, 95% CI: 1.41-13.08) were significantly associated with the outcomes of EDH treatment.

The outcomes of EDH was better than studies conducted in Pakistan 7.9%(12), Nigeria (16.9%)(32) India (13.33%) (9), and Australia (30.8 %). This difference might be due to variation of study population, area and methods used. A study in Australia shows that 21.5% of severe TBI had EDH, while in this study 52(14.2%) of EDH patients were severe TBI. Thus turns to increase the mortality and unfavorable outcomes of EDH.

This study revealed that those who had increased ICP were significantly associated with the poor outcomes (death) of EDH patient. A study in Egypt shows that systolic BP > 90 mmHg at admission significantly associated with favorable outcomes of EDH. Moderate TBI and severe TBI were significantly associated with unfavorable outcomes(33).

This study shows that those EDH patients who had >25mm thickness of hematoma were significantly associated with increased rate death. The reason might be larger hematomas were associated with a higher incidence of progression in size and clinical deterioration causing death.

This study also showed that those patients who had less than 90mmHg systolic blood pressure were significantly associated with the poor outcome (death) of EDH patients.

This study shows that majority of EDH patients were males, 16-30 years and farmers. Fighting and fall down injury were the major causes of EDH. A study in Addis Ababa shows that, assaults and RTAs was the most frequent cause of trauma(31). Also the most common site of EDH was parieto-temporal and frontal region. This finding was similar to a study in Pakistan, temporo-parietal 30.79%, frontal region 26.97%(12), India temporal (20%), and temporo-parietal (20%), parietal (16.66%) and frontal (13.33%)(34).

This study revealed that 57.5% of patients recover well, 37.0% of patients had some disability, 4.3% of patients had severe disability, and 1.2% of patients had persistent vegetative state. This is similar to a study in India, 70% of patients recover well, 20% had moderate disability, 3.33% had severe disability, and 3.33% were in persistent vegetative state(34).

### **6.1 Limitations of the study**

This study was retrospective in nature and is based on the registry book and chart review, which lacks very important variables. This study was institutional based retrospective cross-sectional study and due to the nature of the study design it is not possible to assess the temporal relationship.



## **7. Conclusion and Recommendation**

### **7.1 Conclusions**

The mortality rate of EDH was comparable with other studies. EDH was common in 16-30 years and farmers. The most common presentation of EDH was headache, LOS, vomiting, and increased ICP. The most common site of EDH was Parieto-temporal. Fighting was the leading cause of EDH.

The study identified that increased ICP, >25mm thickness of hematoma and hypotension were significantly associated with the outcomes of EDH treatment.

### **7.2 Recommendation**

**For Minster of health:** Prompt intervention and postoperative management in intensive care definitely improve outcome; create well trained and organized emergency team.

**For health Care Professionals:** should monitor blood pressure and optimize perioperative management of EDH. Prompt and adequate management of EDH shall decrease the mortality of EDH patients. Increased caution and closer monitoring are required when the size of the hematoma is >25 mm.

**For researcher:** I recommend for further study especially using prospective study.

## 8. Reference

1. Khairat A, Waseem M. Epidural hematoma. StatPearls [Internet]: StatPearls Publishing; 2021.
2. Hawryluk GW, Manley GT. Classification of traumatic brain injury: past, present, and future. *Handbook of clinical neurology*. 2015;127:15-21.
3. Lolli V, Pezzullo M, Delpierre I, Sadeghi N. MDCT imaging of traumatic brain injury. *The British journal of radiology*. 2016;89(1061):20150849.
4. Ford LE, McLAURIN RL. Mechanisms of extradural hematomas. *Journal of Neurosurgery*. 1963;20(9):760-9.
5. Eto F, Tatsumura M, Iwabuchi S, Ogawa T, Mammoto T, Hirano A. Clinical features of spontaneous spinal epidural hematoma. *Journal of Rural Medicine*. 2019;14(2):206-10.
6. Bricolo AP, Pasut LM. Extradural hematoma: toward zero mortality: A prospective study. *Neurosurgery*. 1984;14(1):8-12.
7. Narayan RK, Kempisty S. Closed head injury. *Principles of neurosurgery* London: Wolfe. 1994:235-92.
8. El-Fiki A, Halem EA. Conservative Plan for Post Traumatic Extradural Hematoma: Risk Factors Favouring Conversion to Surgery. *Open Journal of Modern Neurosurgery*. 2018;8(3):323-30.
9. Basamh M, Robert A, Lamoureux J, Saluja RS, Marcoux J. Epidural hematoma treated conservatively: when to expect the worst. *Canadian Journal of Neurological Sciences*. 2016;43(1):74-81.
10. Hannah TC, Kellner R, Kellner CP. Minimally invasive intracerebral hemorrhage evacuation techniques: a review. *Diagnostics*. 2021;11(3):576.
11. Balasubramanian H, Saravanan S, Niban G, Raj TG. Efficacy of Decompressive Craniectomy in Acute Subdural Hematoma in Head Injury Patients, Madurai Medical College, Madurai. *INTERNATIONAL JOURNAL OF SCIENTIFIC STUDY*. 2018;6(1):28-32.
12. AYUB S, SHAH M. Management outcome of extradural hematoma. *Pakistan Journal Of Neurological Surgery*. 2014;18(1):17-20.
13. Rosenthal AA, Solomon RJ, Eyerly-Webb SA, Sanchez R, Lee SK, Kiffin C, et al. Traumatic Epidural Hematoma: Patient Characteristics and Management. *The American surgeon*. 2017;83(11):e438-e40.
14. Nasir J, Sheikh F, Benish S, ur Rehman L, Asad A, Ejaz L. Surgical Outcome of Traumatic Posterior Fossa Extradural Hematoma in Paediatric Population: Our Experience at UCHS, The Children's Hospital, Lahore. *Pakistan Journal Of Neurological Surgery*. 2022;26(2):362-9.
15. Biluts H, Abebe M. SHORT-TERM OUTCOME OF OPERATED TRAUMATIC BRAIN INJURY PATIENTS FOR INTRACRANIAL HEMORRHAGE AT TIKUR ANBESSA SPECIALIZED TEACHING HOSPITAL (TASTH), ADDIS ABABA, ETHIOPIA. *Ethiopian Medical Journal*. 2017;55(1):63-8.
16. Bir SC, Maiti TK, Ambekar S, Nanda A. Incidence, hospital costs and in-hospital mortality rates of epidural hematoma in the United States. *Clinical neurology and neurosurgery*. 2015;138:99-103.
17. Meara JG, Greenberg SL. The Lancet Commission on Global Surgery Global surgery 2030: evidence and solutions for achieving health, welfare and economic development. *Surgery*. 2015;157(5):834-5.
18. Macpherson B, MacPherson P, Jennett B. CT evidence of intracranial contusion and haematoma in relation to the presence, site and type of skull fracture. *Clinical radiology*. 1990;42(5):321-6.
19. Rahimi A, Corley JA, Ammar A, Shlobin NA, Rolle M, Mekary RA, et al. The unmet global burden of cranial epidural hematomas: A systematic review and meta-analysis. *Clinical Neurology and Neurosurgery*. 2022;219:107313.
20. Haagsma JA, Graetz N, Bolliger I, Naghavi M, Higashi H, Mullany EC, et al. The global burden of injury: incidence, mortality, disability-adjusted life years and time trends from the Global Burden of Disease study 2013. *Injury prevention*. 2016;22(1):3-18.
21. Irie F, Le Brocque R, Kenardy J, Bellamy N, Tetsworth K, Pollard C. Epidemiology of Traumatic Epidural Hematoma in Young Age. *Journal of Trauma and Acute Care Surgery*. 2011;71(4):847-53.

22. Chicote Álvarez E GCA, Ortiz Lasa M, Jiménez Alfonso A, Escudero Acha P, Rodríguez Borregán JC, Peñasco Martín Y, Dierssen Sotos T. . Epidemiology of traumatic brain injury in the elderly over a 25 year period. *Rev Esp Anestesiol Reanim(Engl ed)*. 2018;65(10):546-51.
23. Sahuquillo J AF. Decompressive craniectomy for the treatment of refractory high intracranial pressure in traumatic brain injury. *Cochrane Database Syst Rev*. 2006(1).
24. Baykaner K AH, Ceviker N, Keskil S, Seckin Z. . Observation of 95 patients with extradural hematoma and review of the literature. *Surg Neurol*. 1988;30:339-41.
25. Zimmerman RA BI. Computed tomographic staging of traumatic epidural bleeding. *Radiology*. 1982;144:809-12.
26. Babu JM PS, Palumbo MA, Daniels AH. Spinal Emergencies in Primary Care Practice. *Am J Med*. 2019;132(3):300-6.
27. Akine Eshete FT. Magnitude of Severe Head Injury and Its Associated Factors among Head Injury Patients in Gedeo Zone, Southern Ethiopia: A Two-Year Retrospective Study. *Ethiop J Health Sci*. 2018;28(3):323.
28. Ayele TA, Zeleke BM, Tessema GA, Melak MF. Magnitude and patterns of injuries among patients in Gondar University Hospital, northwest Ethiopia: an institutional-based study. *Open Access Surgery*. 2017;10:25-31.
29. Walle TA, Tiruneh BT, Bashah DT. Prevalence of head injury and associated factors among trauma patients visiting surgical emergency department of Gondar University Referral Hospital, Northwest Ethiopia 2016. Across-sectional study. *International journal of Africa nursing sciences*. 2018;9:57-61.
30. Assele DD, Lendado TA, Awato MA, Workie SB, Faltamo WF. Incidence and predictors of mortality among patients with head injury admitted to Hawassa University Comprehensive Specialized Hospital, Southern Ethiopia: A retrospective follow-up study. *PloS one*. 2021;16(8):e0254245.
31. Laeke T, Tirsit A, Kassahun A, Sahlu A, Debebe T, Yesehak B, et al. Prospective Study of surgery for traumatic brain injury in Addis Ababa, Ethiopia: trauma causes, injury types, and clinical presentation. *World neurosurgery*. 2021;149:e460-e8.
32. Ayogu OM, Onobun DE, Igbokwe KK, Ugwuanyi CU, Mordi CO, Ibeneme SA. Factors Affecting the Outcome of Traumatic Brain Injured Patients with Acute Epidural Haematoma in National Hospital, Abuja. *Journal of the West African College of Surgeons*. 2021;11(1):1-4.
33. Ssebakumba MK, Lule H, Olweny F, Mabweijano J, Kiryabwire J. Thirty-day clinical outcome of traumatic brain injury patients with acute extradural and subdural hematoma: a cohort study at Mulago National Referral Hospital, Uganda. *Egyptian journal of neurosurgery*. 2020;35(1):1-12.
34. Udaykumar Ramrao Badhe SB. Clinical Evaluation and Management Outcome of Extradural Haematoma. *MVP Journal of Medical Sciences*. 2018;5(1):49–54.
35. Leitgeb J, Mauritz W, Brazinova A, Majdan M, Wilbacher I. Outcome after severe brain trauma associated with epidural hematoma. *Archives of Orthopaedic and Trauma Surgery*. 2013;133(2):199-207.
36. Zwayed ARH, Lucke-Wold B. Conservative management of extradural hematoma: A report of sixty-two cases. *Neurology and clinical neuroscience*. 2018;2(2):5-9.
37. Bedry T, Tadele H. Pattern and outcome of pediatric traumatic brain injury at hawassa university comprehensive specialized hospital, southern Ethiopia: observational cross-sectional study. *Emergency medicine international*. 2020;2020.

## **Annex**

### **Annex I: Information sheet**

Greeting!

My name is Dr. Aytenew Amsalu and I am postgraduate general surgery resident in Bahir Dar University. I'm conducting research on management outcome of epidural hematoma at TGSH and FHCSH, Bahir Dar town, Amhara region, Northwest Ethiopia. The findings of the study will give information about outcome of patients and help conduct further research.

**Title of the research:** management outcome and associated factors of epidural hematoma in FHCSH and TGSH, Northwest Ethiopia 2022

**Objective:** To assess the management outcome and associated factors of epidural hematoma in FHCSH and TGSH, Northwest Ethiopia 2022

**Participants:** Secondary data will be collected from patient chart

For further information: Dr Aytenew Amsalu;

Mob:- +251974497947, e-mail:- [aytenew@gmail.com](mailto:aytenew@gmail.com)

Data collector: \_\_\_\_\_ Sig \_\_\_\_\_

Code: \_\_\_\_\_

**Annex II: English Version Checklist**

Instruction: Please read the questionnaires carefully and circle the alternative and fill the space provided.

***Part I: socio demographic Characteristics***

| s/n  | Questions    | Response  | Skip |
|------|--------------|---|------|
| 101. | Residency    | 1. Rural      2. Urban  |      |
| 102. | Sex          | 1. Male      2. Female  |      |
| 103. | Age in years | _____ years   |      |
| 104. | Occupation   | 1. Farmer      4. Government employee<br>2. Driver      5. Other _____<br>3. Merchant |      |

***Part II: Clinical related characteristics***

|      |   |  |  |
|------|---|--|--|
| 201. | Chronic illness (comorbidity)           | 1. Yes      2. No  |  |
| 202. | If Q201 answer is yes, type of illness? | 1. DM      4. renal disease<br>2. HTN      5. Other _____<br>3. Cardiac diseases |  |
| 203. | History of medication intake            | 1. Anticoagulants      3. CNS drugs<br>2. Antihypertensive      4. Other         |  |
| 204. | History of trauma to the head           | 1. Yes      2. No  |  |

***Part III: Pattern and mechanism of traumatic head injury***

|     |                                  |   |  |
|-----|----------------------------------|---|--|
| 301 | Mechanism of injury              | 1. RTA      4. Animal kick or bite<br>2. Falls      5. Other<br>3. Fighting |  |
| 302 | Time of arrival after injury     | _____hrs  |  |
| 303 | Duration of hospital stay        | _____   |  |
| 304 | Place of occurrence              | 1. Occupant vehicle      3. Outdoor<br>2. Pedestrians      4. Home          |  |
| 305 | History of loss of consciousness | 1. Yes      2. No   |  |

|     |  |  |  |
|-----|--|--|--|
| 306 | If yes to Q. no 305, for how long?               | _____minute  |  |
| 307 | History of abnormal body moment                  | 1. Yes      2. No  |  |
| 308 | History of Headache                              | 1. Yes      2. No  |  |
| 309 | History of Vomiting                              | 1. Yes      2. No  |  |
| 310 | Dominance  | 1. Right-handed      2. Left-handed  |  |
| 311 | GCS  | _____  |  |
| 312 | Loss of consciousness                            | 1. Yes      2. No  |  |
| 313 | Seizure  | 1. Yes      2. No  |  |
| 314 | Aspiration                                       | 1. Yes      2. No  |  |
| 315 | Hypotension (SBP< 90 mmHg                        | 1. Yes      2. No  |  |
| 316 | Increased ICP                                    | 1. Yes      2. No  |  |
| 317 | If yes to Q.no 6, what symptoms of increased ICP | 1. Vomiting      3. Decreased mentation<br>2. Hypertension   |  |
| 318 | Pupillary sign                                   | 1. Midsized & reactive<br>2. Unilaterally dilated &fixed<br>3. Bilaterally dilated &Fixed          |  |
| 319 | Body weakness                                    | 1. No                      3. Hemiparesis<br>2. Monoparesis      4. Hemiplegia                     |  |
| 320 | Severity of TB                                   | 1. Mild                      3. Severe<br>2. Moderate              4. Critical                     |  |
| 321 | Associated other intracranial injury             | 1. Yes                      2. No  |  |
| 322 | If answer to Q. no 11 is yes, which one          | 1. Brain Contusion<br>2. Subdural hematoma<br>3. Intracranial Hemorrhage<br>4. Skull bone fracture |  |
| 323 | Associated Extracranial injury                   | 1. Yes                      2. No  |  |
| 324 | If yes to Q. no 323, which of it                 | 1. Extremity bone fracture<br>2. Chest injury<br>3. Abdominal injury<br>4. Spinal trauma           |  |

|                                  |   |  |  |
|----------------------------------|---|--|--|
|                                  |   | 5. Other (Specify) _____   |  |
| <b>Part IV: CT scan findings</b> |   |  |  |
| 401                              | Sites of EDH  | 1. Frontal      3. parieto-temporal<br>2. Temporal    4) occipital 5) others _____                         |  |
| 402                              | Thickness of the hematoma                             | _____cm  |  |
| 403                              | Involvement of Hemisphere                             | 1. Right      2. Left      3. Bilateral  |  |
| 404                              | Midline shift   | 1. No          2. <5mm      3. >5mm  |  |
| 405                              | 4) Other CT findings                                  | 1. None      2. Linear skull fracture    3. DSF<br>4. Contusion    5. Other(specify) _____                 |  |
| <b>Management given</b>          |   |  |  |
| 406                              | Type of surgical procedure                            | 1. Burr hole      3. Elevation & evacuation<br>2. Craniotomy    4. Others _____                            |  |
| 407                              | Type of anesthesia used                               | 1. Local anesthesia    2. General anesthesia   |  |
| 408                              | ICU admission   | 1. Yes              2. No  |  |
| <b>Management outcome</b>        |   |  |  |
| 409                              | Condition on discharge                                | 1. Improved    2. Death  |  |
| 410                              | If Q410 answer is death, what is the cause of death ? | 1. Primary brain injury    3. Cardiac/Renal failure<br>2. Respiratory failure    4. Others (Specify) _____ |  |
| 410                              | Neurologic outcome                                    | 1. Good              3. Severe disability<br>2. Some disability    4. persistent vegetative state          |  |

**Thank You!**