

2022-08

# Treated Bed Nets Utilization and Associated Factors Among Under Five Children in Lasta District, North Wollo Zone ,Amhara Region, North East Ethiopia,2022

Asfaw, Hailu

---

<http://ir.bdu.edu.et/handle/123456789/15066>

*Downloaded from DSpace Repository, DSpace Institution's institutional repository*



**BAHIR DAR UNIVERSITY**  
**COLLEGE OF MEDICINE AND HEALTH SCIENCES**  
**SCHOOL OF PUBLIC HEALTH**  
**DEPARTMENT OF EPIDEMIOLOGY AND BIOSTATISTICS**  
**INSECTICIDE TREATED BED NETS UTILIZATION AND**  
**ASSOCIATED FACTORS AMONG UNDER FIVE CHILDREN IN**  
**LASTA DISTRICT, NORTH WOLLO ZONE ,AMHARA REGION,**  
**NORTH EAST ETHIOPIA,2022**

**BY: ASFAW HAILU ZERIHUN (FETP RESIDENT)**

**A THESIS SUBMITTED TO THE DEPARTMENT OF EPIDEMIOLOGY AND  
BIOSTATISTICS, SCHOOL OF PUBLIC HEALTH, COLLEGE OF MEDICINE  
AND HEALTH SCIENCES IN PARTIAL FULFILMENT OF THE  
REQUIREMENT FOR THE DEGREE OF MASTERS IN FIELD  
EPIDEMIOLOGY**

**AUGUST, 2022**

**BAHIR DAR**



BAHIR DAR UNIVERSITY  
DEPARTMENT OF EPIDEMIOLOGY AND BIOSTATISTICS,  
SCHOOL OF PUBLIC HEALTH, COLLEGE OF MEDICINE AND  
HEALTH SCIENCES  
INSECTICIDE TREATED BED NETS UTILIZATION AND  
ASSOCIATED FACTORS AMONG UNDER FIVE CHILDREN IN  
LASTA DISTRICT, NORTH WOLLO ZONE, AMHARA REGION,  
ETHIOPIA,2022

BY: ASFAW HAILU (FETP RESIDENT)

ADVISOR

1. KEBADNEW MULAT (MPH, ASSISTANT PROFESSOR)
2. GEBEYAW WUDIE (MPH, ASSISTANT PROFESSOR)

AUGUST, 2022

BAHIR DAR



**Acknowledgment**

I would like to express my gratitude to my academic advisors Mr. Keadnew Mulat and Mr. Gebeyaw Wudie, for their unlimited advice, provision of constructive comments, and guidance that helped me throughout the work to prepare this research thesis. I would also thanks to Bahir Dar university department of Epidemiology and Biostatistics, school of public health, college of medicine and health sciences for providing the opportunity to conduct this public health important title. I would like to thank North Wollo Zone health department, Lasta district health office, and staff for their contribution to the accomplishment of this study. Finally, I would like to thank data collectors and respondents who participate in this study.

## Tables of contents

|   |     |
|---|-----|
| Acknowledgment .....                                | i   |
| List of Table.....                                  | iv  |
| List of figures .....                               | v   |
| Acronyms/Abbreviations .....                        | vi  |
| Abstract .....                                      | vii |
| 1. Introduction.....                                | 8   |
| 1.1. Background.....                                | 8   |
| 1.2. Statement of the problem .....                 | 9   |
| 1.3. Significance of the study.....                 | 10  |
| 2. Literature review .....                          | 11  |
| 2.1. Utilization of ITN in under-five children..... | 11  |
| 2.2. Factors associated with ITN utilization .....  | 11  |
| 2.2.1. Sociodemographic factors.....                | 11  |
| 2.2.2. ITN related factors .....                    | 12  |
| 2.2.3. Knowledge related factors .....              | 13  |
| 2.2.4. Intra-household characteristics.....         | 14  |
| 4. Objective .....                                  | 16  |
| 4.1. General objective .....                        | 16  |
| 4.2. Specific Objectives .....                      | 16  |
| 5. Methods and materials .....                      | 17  |
| 5.1. Study design, area and period .....            | 17  |
| 5.2. Source and study populations .....             | 18  |
| 5.2.1. Source population .....                      | 18  |
| 5.2.2. Study population .....                       | 18  |
| 5.3. Inclusion and Exclusion criteria.....          | 18  |
| 5.4. Sample size determination .....                | 19  |
| 5.5. Sampling technique.....                        | 20  |
| 5.6. Variables of the study .....                   | 21  |

|   |    |
|---|----|
| 5.6.1. Dependent variable.....  | 21 |
| 5.6.2. Independent variables .....  | 21 |
| 5.7. Operational Definitions.....   | 22 |
| 5.8. Data collection methods and procedure .....                            | 22 |
| 5.10. Data processing and analysis .....                                    | 22 |
| 5.12. Dissemination of findings .....                                       | 23 |
| 6. Results.....   | 24 |
| 6.1. Sociodemographic characteristics of care takers .....                  | 24 |
| 6.2. Utilization of ITN in children under five .....                        | 24 |
| 6.2. Intra-household characteristics .....                                  | 25 |
| 6.3. Knowledge related factors .....  | 25 |
| 6.4. Factors associated to utilization of ITN in under- five children ..... | 26 |
| 7. Discussions .....  | 28 |
| 9. Conclusions.....   | 29 |
| 10. Recommendations.....  | 29 |
| 11. References.....   | 30 |
| 12. Annexes.....  | 33 |



### **List of Table**

|  |    |
|--|----|
| Table 1. Sample size determination for associated factors using Epi-info STATCAL: cross-sectional, Flessis for the study done in Last district, Amhara, Ethiopia, 2022 ..... | 20 |
| Table 2. Socio demographic variables of care taker with under-five children in Lasta district, North Wollo Zone, Amhara, Ethiopia, 2022 .....                                | 24 |
| Table 3. Intra-household characteristics in Lasta district, North Wollo zone, Amhara, Ethiopia, 2022.....  | 25 |
| Table 4. Knowledge related factors related to ITN utilization in Lasta district, North Wollo zone, Amhara, Ethiopia, 2022 .....  | 25 |
| Table 5. Logistic regression analysis of variables for utilization of ITN among under-five children in Lasta district, North Wollo, Amhara, Ethiopia, 2022 .....             | 27 |

### **List of figures**

|  |    |
|--|----|
| Figure 1. Conceptual framework adopted from different literature .....   | 16 |
| Figure 2. Map of Lasta district, North Wollo, Amhara region, 2022.....   | 18 |
| Figure 3. Schematic presentation of sampling procedure to select study subjects from Last district, North Wolo zone, Amhara, Ethiopia, 2022..... | 21 |

**Acronyms/Abbreviations**

|       |  |
|-------|--|
| ACT   | Artemisinin-Based Combination Therapy      |
| CI    | Confidence Interval                        |
| ETB   | Ethiopian Birr                             |
| ENMIS | Ethiopia National Malaria Indicator Survey |
| FMHO  | Federal Ministry Health Organization       |
| IEC   | Information Education Communication        |
| IRS   | Indoor Residual Spray                      |
| ITN   | Insecticide Treated Net                    |
| NMSPs | National Malaria Survey Programs           |
| PW    | Pregnant Women                             |
| RBW   | Roll Back Malaria                          |
| SKM   | Square Kilo Meter                          |
| SPSS  | Statistical Package for Social Science     |
| SSA   | Sub-Saharan Africa                         |
| UFC   | Under -Five Children                       |
| WHO   | World Health Organization                  |

## **Abstract**

**Background:** Malaria can be prevented by using insecticide treated nets and decreases malaria mortality by 55% in under -five children in Africa. Ethiopia, scaling up distribution and utilization of insecticide treated nets to cover 100% of under –five children. Utilization of insecticide treated nets has many unsolved problem like, not being hung at all, poorly hung nets, and children not being prioritized to use. This study purpose was to assess insecticide treated nets utilization and its associated factors among under -five children in Lasta district.

**Methods:** we conducted a community-based cross-sectional study among under-five children in Lasta district from April to May, 2022. Four study kebeles were identified by simple random sampling technique and 416 care takers with at least one under five children were selected by systematic random sampling technique. Semi-structured questionnaire was used to collect data using face-to-face interview. The data was entered into Epi data version 3.1 then exported to statistical package for social science version 25 for analysis. Pretest was conducted among 5% of sample. We used tables, frequencies, percentages, and means for descriptive analysis. Bivariable and multivariable logistic regression analysis were done. Variables with p-value <0.05 along with a 95% confidence interval (CI) were used to declared statistically significant variables. Hosmer-Lemeshow model was done.

**Results:** A total of 403 under-five children were participated in the study with 95.72% response rate. Of these, 137 (34%) slept under insecticide treated net during the previous night prior to the survey. Care takers who had social support from others [(AOR: 1.74, 95% CI (1.10-2.74)], who had treated net [(AOR: 3.04, 95% CI (1.07-8.63)], who had good knowledge about insecticide treated net [(AOR: 10.01, 95% CI (5.37-18.69)], Care takers who think the ITN chemical is harmful [(AOR: 0.62, 95% CI (0.39-0.99)] were significantly associated with under five children insecticide treated nets utilization.

**Conclusions:** Even if all have access to insecticide treated nets, only one third of the participant were utilized it. Care takers who had no social support from others, had non treated nets, had poor knowledge about insecticide treated nets and care takers who did not think insecticide treated net chemical is harmful were significantly associated factors. Thus, health education and awareness creation should be taken for those care takers to improve insecticide treated nets utilization for under-five children.

**Keywords:** under-five children, insecticide-treated nets, utilization, malaria, Last district

## **1. Introduction**

### **1.1. Background**

Insecticide-treated bed net (ITN) is a type of vector control approach and cost-effective type of malaria prevention (1). Between 2016 and 2018, a total of 578 million insecticide-treated mosquito nets (ITNs) were reported by manufacturers as having been delivered globally, with 50% going to Côte d'Ivoire, the Democratic Republic of the Congo, Ethiopia, Ghana, India, Nigeria, Uganda and the United Republic of Tanzania. In 2018 about 197 million ITNs were delivered by manufacturers, of which more than 87% were delivered to countries in sub-Saharan Africa (2).

Insecticide treated nets have been proved to reduce all causes child mortality by 14-33 percent in rural sub-Saharan Africa. ITNs have been promoted to protect the most susceptible to severe malaria, children under five (3).

Ethiopia, realizing the effectiveness, scaling up distribution and utilization of ITNs to cover 100% of under-five children (4). ITNs were widely distributed in rural area for free in order to reduce the malaria burden. As a result, the number of new bed owners has increased dramatically. A wide gap exists between coverage and utilization of ITNs in the country (5). Despite the fact that night time usage was best, some people use the net to other purpose (6).

Mosquito net ownership in itself is not synonymous with utilization especially for under-five children (7). Insecticide treated nets are helpful in lowering child mortality, parasite prevalence, and simple and severe malaria episodes (8). Insecticide-treated nets have turned into an important element of malaria prevention since the beginning of the new millennium, known to promote the dramatic decline in disease incidence and malaria-related deaths (9).

In the last decade, the majority of malaria-endemic countries have struggled to build effective national-scale continuous distribution mechanisms for ITNs (10). In Sub-Saharan Africa, half of those at risk of malaria sleep under an ITN; in 2018, 50 % of the population was protected by this intervention, up from 29 % in 2010. Furthermore, from 33% in 2010 to 57 % in 2018, the percentage of the population with access to an ITN increased. Care takers with at least one ITN for every two individuals climbed to 72 % in 2018, from 47 % in 2010. However, this figure indicates just a slight gain over the past 3 years, and remains far from the aim of universal coverage (11).

The use of insecticide-treated nets has been the principal method of malaria control in malaria-endemic areas. Despite its relative success, gains are treated by mosquito resistance to pyrethroids (insecticides used on nets), physical and chemical deterioration of ITNs, which diminishes their efficacy, and inconsistent and inaccurate application by humans, among other problems (12).

Ethiopia's national malaria prevention policy seeks to supply one ITN for every sleeping space (approximately one net for 1.8 persons) (13). To promote universal access to ITNs have adopted policies by most malaria-endemic country (14). Ownership and utilization of mosquito nets for malaria prevention is still sub-optimal with only 62% ownership of at least one mosquito net and only 37% of children using the nets (15).

### **1.2.Statement of the problem**

Malaria is a public health problem with 219 million cases and 435,000 malaria-related deaths globally in 2017(16). It is the number one killer of under-fives children, with one child dying every two minutes (17). Under-five children are the most vulnerable to malaria, accounting for 61 % of all malaria deaths globally(18). In African region malaria burden and deaths from worldwide accounted 88% and 90% respectively. Eighty-eight percent of all deaths in sub-Saharan Africa attributed to malaria occurring in under five children (2).

World Health Organization (WHO) has formulated different strategies to combat the high prevalence of Malaria in the world. The most commonly used methods to prevent mosquito bites are sleeping under an Insecticide Treated Net. Use of ITNs has been shown to reduce malaria case incidence rates by 50% in a range of settings, and to reduce malaria mortality rates by 55% in children aged less than 5 years in sub-Saharan Africa (18)

Despite increasing availability, ITN utilization is still limited in several area and use of ITNs decreases malaria mortality rates by 55% in under-five children in Africa. Most of sub-Saharan Africa countries use of ITNs to prevent mosquito bites. Although increased efforts to improve ITN accessibility, approximately 437,000 children still die from malaria in Africa every year (19).

The burden of malaria, its prevention and control remains a challenge despite the existence of effective technologies (3). Proper use of ITNs can contribute to 90 % of decrease in malaria and 44% of all causes of death in under-five children (20). Burden of malaria is more devastating in

children and causes a considerable amount of workdays economic cost, reduction of productivity and days missed by women and men due to malaria cases of their children (21).

In Ethiopia, approximately 52 million people (68%) live in malaria risk areas, primarily at altitudes below 2,000 meters. Under-five year children are the most vulnerable group affected by malaria(22).According to the 2018 World Malaria Report, after the increased utilization of ITNs, deaths and admissions due to malaria among under-five children decreased by 81% and 73% respectively. Malaria is one of the ten top leading causes of morbidity and mortality among under-five children in Ethiopia (23). Utilization of ITN has many unsolved problem like, not being hung at all, poorly hung nets, and children not being prioritized to use ITN.

A wide gap exists between coverage and utilization of ITNs in the country. This shows that, mosquito net ownership in itself is not synonymous with utilization especially for under-five children. Complaints of nets not being hung at all, poorly hung nets, and children not being prioritized, the problem remains unsolved, and many questions remain unanswered Increase in ITNs access (i.e. household ownership) does not necessarily translate to equal increase in utilization(7).

Under-five children are vulnerable to malaria and targeted as high priority group for ITNs distribution in the country. Therefore, care takers in targeted areas of vulnerable groups have higher chance of getting free of charge ITNs. Despite the rapid and complete coverage of ITNs distribution, reports indicated that still there is poor ITNs utilization (24).

Even though the coverage of ITN in Lasta district was 100%, a total of 1356 malaria cases were reported in 2013 Ethiopia fiscal year. Of this reported cases, 275 (20.3%) were under five children. In this district, ITN utilization for preventing malaria in under five children is not known (25). Therefore, this study was to know the utilization status of Lasta district and its associated factors among under-five children Lasta district, North Wollo Zone, North West Ethiopia.

### **1.3.Significance of the study**

ITN can be one type of cost effective vector control approach for the prevention and elimination of malaria. However, little is known about ITN utilization and associated with the utilization in under- five years old children in the study area yet. Therefore this study finding may serve as preliminary evidence about ITN utilization among under -five children's in the district. This study finding may help to program planners and implementers' about ITN utilization and factors associated to ITN utilization among under- five children. This study finding may also help to the district health office to facilitate elimination program of malaria.

## **2. Literature review**

### **2.1.Utilization of ITN in under-five children**

Worldwide the targets for universal coverage with ITNs aim to achieve at least 80% coverage for ITN ownership and utilization of it (26). A study conducted in African countries showed that the utilization of ITN among under five children was range from 42% to 51 % (27).

A similar study done with Gindiri national malaria indicator survey showed that the percentage of under five children who slept under ITNs have increased steadily and substantially from 6 % to 26.7 % (28).

A study conducted in Sub-Saharan Africa also showed that 55% of the populations were slept under ITN, of which 68% of was under five children (29).

A study done in Congo showed that 68.2% of children slept under ITN the night before the interview (30). In addition, a study in Myanmar showed that 80.5% of under-five children did not sleep under an ITN last night (31).

A study conducted in Malawi showed that the rate for ITN use was 57.8% (95% Confidence interval (CI) 56.1%–59.4%) (19).A study conducted in showed that 591 (58.6%) reported that their under five children slept under the net the night prior to the survey. Less than half (41.2%) slept every night under an LLIN, while 63 (7.5%) caregivers mentioned their under -five children had never slept under the net(7).

### **2.2.Factors associated with ITN utilization**

#### **2.2.1. Sociodemographic factors**

In a study conducted in western Kenya to assess factors affecting ITNs use the variables that had significant effect on ITNs use was age, for which a 14.5% reduction in the probability of uses was observed in children under five years of age as compared to individuals' greater than five years of age (32).

A study conducted in Ethiopia ITNs use was more common when the house is headed by a female (OR=1.92, 95% CI = 1.34 - 2.74). Some 65 (17.06%) respondents were from female-headed households. Female respondents constituted 319 (83.73%). Women who believe that it is normal to use ITNs were 1.9 times more likely to use ITN than those who did not (OR: 1.930; 95% CI: 1.645–2.265).On the other hand, that ITNs use was nearly similar for under-five children of both genders, for boys it was 65.4% and for girls 64.1%). Similarly this found that there was no significant association on gender of household head and utilization of ITNs by under-five children(22).



A study showed that in Ghana the most (92.6%) of the respondents was Christians and Similar study in Myanmar showed that one hundred and three respondents (64.4%) were Christians than Muslim(31).

Care takers that had two or three children also had a great likelihood of bed net use than in the households with four or more children (33).

Education in Ethiopia showed that a determinant of mosquito net use among vulnerable group. It leads to lack of utilization of the ITN by under-five children in Ethiopia. Children whose mothers/caretakers were literate had 59.4% less risk of malaria compared to those who had not literate(31).A study conducted in Mugisha showed that possession and use of ITNs in children with a married mother/caretakers are over 3 times as likely to sleep under a net (33).

A study conducted in Rwanda children born to a married mother/caretakers or mother living with a partner were 57% less likely (0.43 [0.36–0.52]) to sleep under a bed net compared to children born to a widowed, divorced mother (34).A study done in Ghana showed that Employed caregivers were also 1.51 times more likely to use an ITN than unemployed caregivers and there was no association (AOR=1.51, 95%CI=0.47, 4.92, p=0.490) (27)

### **2.2.2. ITN related factors**

A study done in southeast Asia showed that the information on the utilization of ITNs, treated or untreated, is beneficial for investigating factors associated with their use, ITN are according to international targets (35).

The source of ITN for under five children was distributed by various projects implemented within the health facility catchment area, and donation by non-governmental organizations 9(2.36%) (36).

A study done in southeast Asia showed that the information on the utilization of ITNs, treated or untreated, is beneficial for investigating factors associated with their use, ITN are according to international targets (35).

A study conducted in Sub- Saharan Africa showed that high ITN coverage rates following free bed nets distribution were previously reported in the Congo and elsewhere including Sierra Leone (87.6%), Ethiopia (91.0%), and Togo (96.7%). The majority of 30 (60%) of respondents have ITN while 20 (40%) of respondents have not ITN all households (37).In a study done in Ghana showed that the majority (80.6%) owned ITNs and 41.7% of them slept under the ITNs with their children the night before the survey one(5).

A study conducted in Amhara and Oromia regions shows that ‘conical’ shaped nets were more preferred than ‘rectangular’ shaped ones (OR=2.27 (95% C.I=1.10-4.68). These studies showed that type of ITN are important determinants for prevention and control of malaria among under-five children (38). A study showed in Shewa Robit town 20% of study participants said that using mosquito net was not comfortable because of becoming hot during sleeping and its bad smell. This shows the fear of discomfort may hinder utilization of mosquito nets by under-five children (8).

### **2.2.3. Knowledge related factors**

A study conducted in Ghana showed that lack of access to ITNs and poor knowledge and perception on ITNs and malaria have been previously reported as important barriers to the use of ITNs in parts of Africa. However, access does not always result in usage due to sociocultural and logistical reasons. Evidence from some parts of Ghana has shown that over 40% of ITNs available in the households go unused (27).

A study done in Western Kenya showed that a wide gap exists between coverage and utilization of ITNs in the country. This shows that, mosquito net ownership in itself is not synonymous with utilization especially for under-five children (34).

A study done in West Gojjam zone , Mihrab Abaya Gamo Gofa Zone and Itang, Gambella zone which showed that 72% have poor knowledge to use ITN (AOR=0.28, 95% CI 0.15-0.53) among caregivers aged 36 and above, 82.7% owned an ITN and 39.4% of them slept in it the night before the survey (18).

A study conducted in Nigeria showed that chemicals used to treat the nets were thought to be harmful to adults, children and pregnant women especially affecting breathing a fact that is also reported here as suffocation. As expected, adult child caregivers make decisions to protect their children from adverse reactions emanating from the use of ITNs (39).

Study conducted in East belesa District showed that 202 (27.9%) of the respondents had low levels of knowledge. A majority (82%) of the respondents knew that mosquito bites are causing malaria(40).

According to the study in sierra Leon that approximately 10.0% of caregivers didn’t know that malaria parasites can be transmitted through mosquito bites(41).A similar study found in Ghana a statistically significant relationship between ITN knowledge and use ( $P=0.002$ ; AOR=1.39, 95% CI=0.25-7.56). Those who were knowledgeable about ITNs were found to be more likely to

use them than those who were not(42). Knowledge of malaria in a large sample of residents in Zambia and Zimbabwe was good, and knowledge of the protective efficacy of ITNs was associated with bed net use (43).

On the contrary a study in Afar Region identified that women who got information about malaria and ITNs had positive attitude towards ITNs than those who had not received mothers/care takers (AOR=0.406: 95% CI 0.26-0.63). knowledge of the cause of malaria were associated with the use of ITNs, but the association was border line (44). A study done in Moqudisho showed that the majority of the respondents believed there is a clear benefit of ITNs uses shows that 46 (92%) said yes, 2 (4%) said no and 2 (4%) said don't know (37).

A Study conducted in Nigeria showed that care givers who had good knowledge 476(62.1%) were 1.8 times more likely (AOR= 1.8, 95% CI; 1.4 – 2.5) to utilize ITN for under five year children (35).

#### **2.2.4. Intra-household characteristics**


There are factors that determine net ownership and usage by members sleeping within the household. These factors affect effective utilization of ITNs by household members including children under five. In the larger sample, the addition of one household member reduces the likelihood of a child sleeping under a net by 17%. This is contrary to the results in which possession was the dependent variable (45).

A study in south west Nigeria showed that Children from households with 1-3 members (AOR= 2.29, CI= 1.45-3.63, p= 0.001) and 4-6 members (AOR= 1.50, CI= 1.04-2.15, p= 0.03) were more likely to sleep under an ITN than children from households with 7 and above members. Care takers that had two or three children also had a great likelihood of bed net use than in the households with four or more children (33) Children from Osun (AOR= 0.49, CI=0.27-0.87, p= 0.016), Ekiti (AOR= 0.34, CI= 0.19-0.60, p= 0.001) and Lagos states (AOR= 0.49, CI= 0.26-0.90, p= 0.022) were less likely to sleep under an ITN than those from Ogun Stateew (46).

A study done in Mpanda showed that Children under five use of any net by children under five in Mpanda was higher (84%) compared to (56%) in Kisarawe (OR: 6.25, p < 0.001) (32). According to a study conducted in Sub-Saharan Africa, by 2019, 68 % of households had at least one ITN. The percentage of the population sleeping under an ITN increased significantly for the whole population (from 2% to 46%) and for children under -five (from 3% to 52%) (47). A study done in Tanzania showed that an increase of 10 % ownership of at least one mosquito net at

household level is associated with 12 % and 10 % decrease of malaria risk for all age and under-five children per year, respectively (48).

### 3. Conceptual framework






Figure 1. Conceptual framework adopted from different literature

## 4. Objective

### 4.1. General objective

- To assess the utilization of ITNs and its associated factors among under-five children in Lasta district, North Wollo zone, Amhara region, Ethiopia April to May, 2022.

### 4.2. Specific Objectives

- To determine the level of utilization of ITNs among under five children in Lasta district, North Wollo zone, Amhara region, Ethiopia April to May, 2022.
- To identify factors affecting utilization of ITN among under-five children in Lasta district, North Wollo zone, Amhara region, Ethiopia April to May, 2022.

## **5. Methods and materials**

### **5.1. Study design, area and period**

A community-based cross-sectional study was conducted from April 1 to May 30, 2022 in Lasta district. Lasta district is one of the districts in North Wollo zone of the Amhara region, Northern Ethiopia. Lalibella is a town of Lasta district. The town is situated 701 km away from Addis Ababa, 305 km away from regional town Bahir Dar, and 180 km away from Zonal town Woldia. Lasta district is bounded; on the South by Meket district, on the West by Bugna district, on the North by Waghumra special zone, and on the East by Gidan district. The district has a total of 26 kebeles. The rainfall ranges between 4200 to 1600 millimeters per year. The total population of district is 142,965. Of which 90,897 (63.58 %) population is living in malaria-risk areas. There are 8 health centers, 27 health posts, and 1 private clinic in the district.




Figure 2. Map of Lasta district, North Wollo, Amhara region, 2022

## 5.2. Source and study populations

### 5.2.1. Source population

All under-five children who have resided in all kebeles of Lasta district were the source population

### 5.2.2. Study population

All under-five children who have resided in the four selected kebeles of Lasta district were the study population

## 5.3. Inclusion and Exclusion criteria

In this study, under-five children who have resided in the selected kebeles of the Lasta district were included. Besides, those under-five children who had non-resident care takers (i.e. who have resided for < 6 months) were excluded as people who have resided for < 6 months in a given administrative unit are not officially recognized as residents.

#### 5.4. Sample size determination

**For specific objective 1:** The sample size was determined using single population proportion formula for the proportion (p) of ITN utilization of children slept under ITN in the previous night prior to the survey in East Belesa district, Amhara region (56.5%) taken from a previous study (40). Assuming, level of significance 95%, 5% margin of error. The sample size of the study was computed using Epi-info Version 7.1.

$$n = Z^2 p (1-p) / d^2, \text{ Where } Z=1.96 \quad p=0.57 \quad d=0.05$$

$$= (1.96)^2 * 0.57(1-0.57) / (0.05)^2$$

$$= (0.9416) / 0.0025 = \underline{\underline{378}}$$

The final sample size with 10% non-response rate was 416.

**For specific objective 2:** Sample size was calculated for the associated factors based on the following assumptions. A 95% level of confidence, 80% of power to detect real association of exposure variable and exposed to unexposed ratio each variable. To determine the sample size frequency of exposure variables Age, Education, Family sizes were taken from different literature. Epi info software was used to determine sample size of each factor by using each odds ratio. We were taking the factors that have an association with ITN utilization from previous studies done in Ethiopia. The factors were summarizing by the following table.



Table 1. Sample size determination for associated factors using Epi-info STATCAL: cross-sectional, Flessis for the study done in Lasta district, Amhara, Ethiopia, 2022

| Variable    | Categories   | Variable (exposed) | Variable (unexposed) | AOR   | % of ITN Utilization among exposed | % of ITN utilization Among unexposed | Calculate Sample size | Non-response rate (10%) | Total sample size | Reference |
|-------------|--------------|--------------------|----------------------|-------|------------------------------------|--------------------------------------|-----------------------|-------------------------|-------------------|-----------|
| Age         | 31–44years   | 11                 | 217                  | 0.05  | (40%)                              | (5%)                                 | 151                   | 161                     | 161               | (24)      |
| Family size | ≤5 families  | 122                | 41                   | 11.23 | (75 % )                            | (6%)                                 | 59                    | 65                      | 65                | (24)      |
| Education   | No education | 200                | 575                  | 3.1   | (25%)                              | (26%)                                | 132                   | 145                     | 145               | (29)      |

### 5.5. Sampling technique

Among the nineteen malarious kebeles identified, 4(20%) of the malarious kebeles was selected randomly by lottery simple random sampling method. The total number of under-five children with care taker in each kebele was obtained using the family folder registration as a sampling frame in the community health information system of the kebeles in the health post. We used the proportional allocation to select study participants from selected kebeles. To select each study unit systematic random sampling method was used. The  $K^{\text{th}}$  were calculated for Kebele Gelesot, Erfa, Dibko and Ayedefer were 4, 6, 4, and 3 respectively. The first study participant was selected randomly from each Kebele.




Figure 3. Schematic presentation of sampling procedure to select study subjects from Last district, North Wollo zone, Amhara, Ethiopia, 2022.

## 5.6. Variables of the study

### 5.6.1. Dependent variable

- The outcome variable was ITN utilization of under- five children (utilized, not utilized).

### 5.6.2. Independent variables

#### Socio-demographic characteristics:

- Care taker age, sex
- Education,
- Occupation, marital status and religion

**Knowledge related factors:** Knowledge of malaria, malaria symptoms, transmissions, prevention, and ITN prevent from malaria

#### ITN related variables:

- Type of ITN, source of ITN, and number of ITN observed, fear of side effect.

## 5.7. Operational Definitions

**ITN utilization :** A child who was reported to have slept under ITNs during the night prior to the survey date (24).

**Caretaker:** - A family member who regularly looks after a child under the age of five.

**Good knowledge:-** of the seven asked knowledge questions, who answered five and above questions correctly (scored 70% and above) (49).

**Poor knowledge:-** those who scored below 70% of the score value of the knowledge measuring question (49).

## 5.8. Data collection methods and procedure

A semi-structured questionnaire was used for data collection. Data were collected by a combination of face-to-face interview and observation. The data collectors and supervisors were trained for two days by the principal investigator. We recruited four diploma nurses as a data collector and two health officers as supervisors. ITN use was cross checked with the interview response of caretakers, through direct observation. The supervisors and principal investigator were assigned to supervise the data-collection process.

Data collection tools contain information on the socio-demographics factors, intra-household factors, Knowledge related factors, and ITN related variables.

## 5.9. Data Quality control

The questionnaire was prepared originally in English and translated to Amharic and back to English to check the quality of the questions. Pre-test was conducted the total sample size to ensure that the questionnaires were free from ambiguity and that the data generated was meaningfully analyzed. It was done by administering the questionnaire in one of neighboring Kebele named Shallo, located 35 Kms from the study area. The principal investigator supervised the overall data collection process. On a daily basis, both data collectors and supervisors were received feedback on the previous day's operations.

## 5.10. Data processing and analysis

Data were cleaned, coded, and entered into Epi data version 3.1 then export to SPSS version 25 for statistical analysis. Descriptive statistics were used to summarize and present the socio-demographic, intra-household factors, Knowledge factors, ITN related variables. The results were presented by text and in the form of tables using frequencies and percentages. Bivariable logistic regression analysis was used to identify factors which has a p- value of less than 0.25 and

considered for multivariable logistic regression analysis. A p-value of less than 0.05 with a 95% confidence interval in multivariable logistic regression analysis was considered as statistically significant factors. The Hosmer–Lemeshow goodness-of-fit was used to test model fitness (With a p-value of 0.06).

#### **5.11. Ethical consideration**

Ethical clearance was obtained from the Institutional Review Board of Bahir Dar University's College of Medicine and Health Science's school of public health department of Epidemiology and Biostatistics. Support letter was obtained from North Wollo zone, Lasta district health office, and from selected kebeles authority. Verbal consent was obtained from the participants care taker. Participants had the right to refused to participate in this research and they have been full right to withdraw from the interview at any time they wish. Finally, participant's identity was kept unknown throughout the data collection and analysis process.

#### **5.12. Dissemination of findings**

The finding of this study was submitted to the department of public health, Bahir Dar University college of Medicine and Health Sciences School of Public Health Department of Epidemiology and Biostatistics and North Wollo zone health department, Lasta district health offices and administrators at Lasta district.

## 6. Results

### 6.1. Sociodemographic characteristics of care takers

A total of 403 caretakers with under-five children were participated in the study with 95.72% response rate. Three hundred two (75%) of the caretakers were females. The mean age of the caretakers was 35 years. A total of 229 (56.8%) participants were unable to read and write (details shown below in Table. 2).

Table 2. Socio demographic variables of care taker with under-five children in Lasta district, North Wollo Zone, Amhara, Ethiopia, 2022

| Variables                        | Category                 | Frequency (%) |
|----------------------------------|--------------------------|---------------|
| Sex of care taker                | Female                   | 302(75)       |
|                                  | Male                     | 101(25)       |
| Age of caretaker                 | 18-24                    | 7(2)          |
|                                  | 25-32                    | 180(45)       |
|                                  | >32                      | 216(53)       |
| Educational status of care taker | Unable to read and write | 229(56.8)     |
|                                  | able to read and write   | 118(29.3)     |
|                                  | Primary                  | 20 (5.0)      |
|                                  | Secondary                | 23 (5.7)      |
|                                  | Above secondary          | 13 (3.2)      |
| Religion                         | Orthodox                 | 364 (90.3)    |
|                                  | Muslim                   | 39 (9.7)      |
| Marital status                   | Married                  | 352 (87.0)    |
|                                  | Divorced                 | 35 (9.0)      |
|                                  | Widowed                  | 11 (3.0)      |
|                                  | Single                   | 5 (1.0)       |
| Occupation of care taker         | Farmer                   | 276 (68.5)    |
|                                  | House Wife               | 103 (25.6)    |
|                                  | Student                  | 7 (1.7)       |
|                                  | Unemployed               | 9 (2.2)       |
|                                  | Daily Laborer            | 8 (2.0)       |

### 6.2. Utilization of ITN in children under five

A total of 137(34%) at 95% CI (29-39.2) under five children were slept under ITN during the night preceding the survey. The number of ITNs of reported 403, observed 304 by data collectors and hanged over the bed 137 had great variation in number.

## 6.2. Intra-household characteristics

A total of 81 (20.0%) care takers had only one separate bed room. About 382 (86.3%) of care takers had two or more sleeping spaces. Four hundred three (100%) of the caretakers had ITN in their house (details shown below in the table 3).

Table 3. Intra-household characteristics in Lasta district, North Wolo zone, Amhara, Ethiopia, 2022.

| Variables                     | Categories      | Frequency (%) |
|-------------------------------|-----------------|---------------|
| Number of under five children | 1               | 185 (45.9)    |
|                               | $\geq 2$        | 223 (55.3)    |
| Family size                   | 1-4             | 190 (46.0)    |
|                               | 5-7             | 195 (48.0)    |
|                               | $\geq 8$        | 19 (0.02)     |
| Number of sleeping space      | One             | 81 (20.0)     |
|                               | Two             | 287 (71.0)    |
|                               | Three and above | 35 (9.0)      |
| Number of ITN in the House    | 1               | 198 (49.1)    |
|                               | 2               | 187 (46.4)    |
|                               | $\geq 3$        | 18 (4.5)      |
| Social support to under five  | Yes             | 166 (41.2)    |
|                               | No              | 237 (58.8)    |

## 6.3. Knowledge related factors

Overall, a total of 280 (70.0%) under five children's care taker had good knowledge related to ITN utilization. Care takers who had known ITN used prevent from malaria were 108 (26.8%). One hundred thirty seven (34%) of under-five children's care takers had ever heard about ITN (details shown below in the table 4).

Table 4. Knowledge related factors related to ITN utilization in Lasta district, North Wollo zone, Amhara, Ethiopia, 2022

| Variables                       | Frequency (%) |
|---------------------------------|---------------|
| Knowledge about ITN utilization | 280 (70.0%)   |
| Knowledge of malaria            | 403 (100%)    |
| Malaria symptoms                | 360 (89.3%)   |
| Transmissions                   | 108 (26.5%)   |
| Prevention                      | 403 (100%)    |
| Ever heard about ITN            | 137 (34%)     |
| ITN prevent from malaria        | 108 (26.8%)   |

#### **6.4. Factors associated to utilization of ITN in under- five children**

For the analysis of bivariable logistic regression, twelve variables to ITN utilization among under-five children were considered. Variables with a p-value of less than 0.25 in the bivariable logistic regression analysis were nominated for the final multivariable logistic regression analysis. For the analysis of multivariable logistic regression seven variables were considered and analyzed.

The odds of care takers who had social support were 2 times more likely (AOR: 1.74, 95% CI (1.10-2.74)) utilized the ITNs for under-five children as compared to those who had not social support. The study also showed that the odds of utilization of ITN was 3.04 times more likely higher among care takers who had treated ITN than who had non- treated ITN (AOR: 3.04, 95% CI (1.07-8.63)).

The odds of care takers who had good knowledge were 10 times more likely utilized ITNs for their children as compared to who had poor good knowledge (AOR: 10.01, 95% CI (5.37-18.69)). The odds of care takers who think the ITN chemical is not harmful were 38% less likely utilized for their children as compared to than who think the ITN chemical is harmful (AOR: 0.62, 95% CI (0.39-0.99)).

Table 5. Bivariable and multivariable logistic regression analysis of variables for utilization of ITN among under-five children in Lasta district, North Wollo, Amhara, Ethiopia, 2022.

| Variables  | Utilization of ITN |              | Bivariable analysis | Multivariable analysis |                     |
|--|--------------------|--------------|---------------------|------------------------|---------------------|
|  | utilized           | Non-utilized | COR (95% CI)        | AOR (95% CI)           | P-value             |
| <b>Sex of care taker</b>                               |                    |              |                     |                        |                     |
| Male   | 68 (25.6%)         | 33 (24.1%)   | 1.08 (0.57-1.49)    |                        |                     |
| Female   | 198 (74.4%)        | 104 (75.9%)  | 1                   |                        |                     |
| <b>Religion of care takers</b>                         |                    |              |                     |                        |                     |
| Orthodox   | 23 (8.6%)          | 16 (11.7%)   | 0.72 (0.37-1.41)    |                        |                     |
| Muslim   | 243 (91.4%)        | 121(88.3%)   | 1                   |                        |                     |
| <b>Partner discuss utilization ITN</b>                 |                    |              |                     |                        |                     |
| Yes  | 210 (78.9%)        | 90 (65.7%)   | 1.96 (1.24-3.10)    | 2.12(1.19-3.81)        | 0.37                |
| No   | 56 (21.1%)         | 47 (34.3%)   | 1                   | 1                      |                     |
| <b>Ever talked importance of sleeping under an ITN</b> |                    |              |                     |                        |                     |
| Yes  | 65 (24.4%)         | 54 (39.4%)   | 0.53(1.29-3.13)     | 0.77(0.43-1.39)        | 0.38                |
| No   | 201(75.6%)         | 83 (60.6%)   | 1                   | 1                      |                     |
| <b>Social support from others</b>                      |                    |              |                     |                        |                     |
| Yes  | 74 (54.0%)         | 92 (34.6%)   | 2.22 (1.46-3.38)    | 1.74(1.10-2.74)        | 0.04 <sup>¥</sup>   |
| No   | 63 (46%)           | 174 (65.4%)  | 1                   | 1                      |                     |
| <b>Insecticide used in ITN is harmful</b>              |                    |              |                     |                        |                     |
| Yes  | 107 (40.2%)        | 80 (58.4%)   | 0.48 (0.32-0.73)    | 0.62 (0.39-0.99)       | 0.0001 <sup>¥</sup> |
| No   | 159 (59.8%)        | 57 (41.6%)   | 1                   | 1                      |                     |
| <b>ITN hanged properly</b>                             |                    |              |                     |                        |                     |
| Yes  | 51(37.2%)          | 67 (25.2%)   | 1.76 (1.13-2.74)    | 1.27(0.72-2.24)        | 0.42                |
| No   | 86 (62.8%)         | 199 (74.8%)  | 1                   | 1                      |                     |
| <b>What type of ITN do you have</b>                    |                    |              |                     |                        |                     |
| Treated  | 259 (97.4%)        | 114 (83.2%)  | 7.47 (3.11-17.89)   | 3.04 (1.07-8.63)       | 0.0001 <sup>¥</sup> |
| Non-treated  | 7 (2.6%)           | 23 (16.8%)   | 1                   | 1                      |                     |
| <b>Knowledge about ITN</b>                             |                    |              |                     |                        |                     |
| Good   | 121(88.3%)         | 124 (46.6%)  | 8.66 (4.88-15.38)   | 10.01(5.37-18.69)      | 0.0001 <sup>¥</sup> |
| Poor   | 16 (11.7%)         | 142 (53.4%)  | 1                   | 1                      |                     |

¥p-value < 0.05 were declared statistically significant variables



## 7. Discussions

This study was conducted to determine ITNs utilization and its associated factors among under-five children of Lasta district. In this study a total of ITN utilization among study subjects in the district was 34 %. This study was lower than the studies done in Sub-Saharan Africa slept under ITNs which is 68 % of them being children under the age of five (29). A study done in Congo showed that the caretakers' of children under five reported that 68.2% of children slept under ITN the night before the interview (30). This finding also different from the study conducted in Malawi which had the rate for ITN use was (AOR:57.8% ,95 %CI (56.1%–59.4%). This difference might be due malaria prevention or elimination strategic activities based on context of the country, geographical and demographic settlement of the community, climatically variations. The odds of care takers who responded that insecticide used in ITN was harmful for under-fives children were 38% less likely (AOR=0.62, 95%CI: 0.39-0.99) not to utilize an ITN as compared to those who said insecticide used in ITN was not harmful. Study conducted in India Ho municipality. The odds of caretakers who responded that ITNs have side effects were 49% times less likely (AOR:0.37, 95%CI (0.22- 0.61) to use an ITN as compared to those who responded had said not harmful (50). This difference might be due to gap of knowledge about ITN, educational status of the caretakers.

There was an association between ITN use and social support for children under five. The odds of care takers who had social support for children under the age of five were 2 times more likely(AOR:1.74,95CI (1.10-2.74) to utilize an ITN compared to those who said that social support had not influence ITN utilization among children under the age of five. This might imply that caregivers had good awareness about ITN utilization for under- five year children(51). The ITN utilization among children under five caretakers who did use treated ITN were 3 times more likely (AOR: 3.04 95%CI: 1.07-8.63) to utilize ITN for their children. This might imply they may think treated ITN is effective to prevent mosquito bite, due to effect of its chemical relative to non -treated ITN.

This study showed that the overall good level of knowledge among care takers was 245(60.8%), whereas 158(39.2%) had poor knowledge about ITN utilization. The good knowledge of caretakers were 10 times more likely (AOR:10.01, 95%CI:5.37-18.69) to utilize ITN for their children when compared to study conducted in Nigeria which showed that 476(62.1%) of care takers had good knowledge about ITN utilization(35).The care takers had nearly similar

knowledge level about ITN utilization. This might be due to usage of similar method of study design and variables used. When compared to study conducted in East belesa District showed that 202 (27.9%) of the respondents had low levels of knowledge(40). This discrepancy might be due to the difference in variables, climatically condition and other geographical reasons.

## **8. Limitations of the study**

This was a cross-sectional study, representing the snapshot of the population within the study period and does not show cause and effect since the independent and outcome variables were measured simultaneously. Data was collected through reporting and thus there was a possibility of recall bias. In this study, however, respondents were required to only recall care taker with under -five children slept under an ITN in the previous night.

## **9. Conclusions**

In the study, the overall utilization of insecticide treated nets by under-fives was low despite the high proportion availability of ITN within the Households with under five children. Besides, even some households with ITN weren't using them consistently. Care takers who had no social support from others, had non treated nets, had poor knowledge about insecticide treated nets and care takers who did not think insecticide treated net chemical is harmful were significantly associated factors.

## **10. Recommendations**

Based on the findings of the study we recommended to:

- Lasta district health office with the respective health care providers should improve ITN utilization to under-five children in the district
- The health care professionals such as health extension workers and health professionals should improve the knowledge of care takers related to ITN utilization
- The family should strengthen the social support of care takers for their children to increase utilizing of the ITN
- Health workers should give awareness to those care takers as insecticide chemical is not harmful for the health of under five children

## 11. References

1. Audu O, Anefu G, Ukpabi D, Anja D, Anejo-Okopi J. The Use of Insecticide-Treated Bed Nets for Control of Malaria in Benue State, Nigeria: The Gap between the Gown and the Town. *JOURNAL OF RESEARCH IN BASIC AND CLINICAL SCIENCES*. 2019;1(1):47-52.
2. Organization WH. The "World malaria report 2019" at a glance. World Health Organization, Geneva, Switzerland. 2019.
3. Sina OJ. An Assessment of the Use of Insecticide Treated Net among children in Ekiti State as a Means of Prevention of Malaria.
4. Admassie A. Insecticide treated nets and associated factors among under five years old children in Mirab -Abaya District,Gamo-Gofa. 2018.
5. Ahorlu CS, Adongo P, Koenker H, Zigirumugabe S, Sika-Bright S, Koka E, et al. Understanding the gap between access and use: a qualitative study on barriers and facilitators to insecticide-treated net use in Ghana. *Malaria journal*. 2019;18(1):1-13.
6. Melaku Birhanu Alemu<sup>1\*</sup> MAA, Melaku Yilma Lemma<sup>3</sup>, Melkitu Fentie Melak<sup>4</sup>, Yenit<sup>5</sup> aMK. Utilization of insecticide treated bed net and associated factors among households of Kola Diba town, North Gondar, Amhara region, Ethiopia. 2018.
7. Lencha B. Insecticide treated net utilization among under five children and household net ownership in Adami Tullu district, Oromia Regional State, Ethiopia: A community based cross-sectional study. *Qual Primary Care*. 2017;26:38-44.
8. Azeb SR, Legesse TW, Hailu FD. Proper utilization of long lasting treated net and associated factors at Zuway Dugda district, Arsi zone, Ethiopia. *Journal of Public Health and Epidemiology*. 2019;11(1):13-24.
9. Pryce J, Richardson M, Lengeler C. Insecticide-treated nets for preventing malaria. *Cochrane Database of Systematic Reviews*. 2018(11).
10. Yukich J, Stuck L, Scates S, Wisniewski J, Chacky F, Festo C, et al. Sustaining LLIN coverage with continuous distribution: the school net programme in Tanzania. *Malar J*. 2020;19(1):158.
11. Giersing BK, Vekemans J, Nava S, Kaslow DC, Moorthy V, Committee WHOPdfVA. Report from the World Health Organization's third Product Development for Vaccines Advisory Committee (PDVAC) meeting, Geneva, 8-10th June 2016. *Vaccine*. 2019;37(50):7315-27.
12. Ngonghala CN, Wairimu J, Adamski J, Desai H. Impact of adaptive mosquito behavior and insecticide-treated nets on malaria prevalence. *Journal of Biological Systems*. 2020;28(02):515-42.
13. UNICEF. Long Lasting Insecticidal Nets Supply Update, New York.
14. Hamre KE, Ayodo G, Hodges JS, John CC. A mass insecticide-treated bed net distribution campaign reduced malaria risk on an individual but not population level in a highland epidemic-prone area of Kenya. *The American journal of tropical medicine and hygiene*. 2020;103(6):2183-8.
15. Idoko L, Okafor,, K.C. A, G.M., Idika, C.N. and Oguiche,, B.E. Assessment of the Willingness to Buy and the use of Insecticide Treated Nets (ITN) among Caregivers of Under-Five Children Attending Immunization Clinics in a Private Tertiary Hospital in Plateau State, Nigeria. 2020.
16. Dhiman S. Are malaria elimination efforts on right track? An analysis of gains achieved and challenges ahead. *Infectious diseases of poverty*. 2019;8(1):1-19.
17. Nnamonu EI, Ndukwe-Ani PA, Imakwu CA, Okenyi CI, Ugwu FJ, Aniekwe MI, et al. Malaria: Trend of Burden and Impact of Control Strategies. *International Journal of TROPICAL DISEASE & Health*. 2020:18-30.
18. Dechasa Nesga<sup>1</sup> DA, Girma Birhanu<sup>1</sup>, Addamu Addissie. Assessment of Malaria Vector Control Measures (ITNs & IRS) Utilization and Factors Affecting it in Adama District, East Shoa Zone, Oromia Region Ethiopia. 2018.

19. Nkoka O, Chipeta MS, Chuang Y-C, Fergus D, Chuang K-Y. A comparative study of the prevalence of and factors associated with insecticide-treated nets usage among children under 5 years of age in households that already own nets in Malawi. *Malaria journal*. 2019;18(1):1-10.
20. Esuabom M, Aworh MK, Umeokonkwo CD, Balogun MS, Bamgboye E, Dahiru T. Utilization of Insecticide Treated Nets and Health Seeking Behavior for fever among Children under Five in Southwest Nigeria. 2021.
21. Ahmed A, Mulatu K, Elfu B. Prevalence of malaria and associated factors among under-five children in Sherkole refugee camp, Benishangul-Gumuz region, Ethiopia. A cross-sectional study. *PLoS one*. 2021;16(2):e0246895.
22. Mulugeta T, Ashenefe B, Shimeka A, Abate S. Insecticide treated net utilization and associated factors among households having under-five children and pregnant women in Fogera District, Northwest Ethiopia. *Ethiopian Journal of public health and nutrition*. 2020;2(2).
23. Organization WH. World malaria report 2020: 20 years of global progress and challenges. *World malaria report 2020: 20 years of global progress and challenges* 2020.
24. Admasie A, Zemba A, Paulos W. Insecticide-treated nets utilization and associated factors among under-5 years old children in Mirab-Abaya District, Gamo-Gofa Zone, Ethiopia. *Frontiers in Public Health*. 2018;6:7.
25. Lasta District Health Office Annual Report. Annual Report,. 2013 E.C.
26. Scott J, Kanyangarara M, Nhama A, Macete E, Moss WJ, Saute F. Factors associated with use of insecticide-treated net for malaria prevention in Manica District, Mozambique: a community-based cross-sectional survey. *Malaria journal*. 2021;20(1):1-9.
27. al. KDKe. Utilization of Insecticide Treated Bed Nets (ITNs) among Caregivers of Children under Five Years in the Ho Municipality. Published 1 April 2019.
28. Sodipo O, Gwomson D, Chingle M, Bupwatda W. The risk factors for malaria in under-fives in Gindiri community of Mangu Local Government Area, Plateau State. *Jos Journal of Medicine*. 2019;13(1):2-12.
29. Zerdo Z, Bastiaens H, Anthierens S, Massebo F, Masne M, Biresaw G, et al. Long-lasting insecticide-treated bed net ownership, utilization and associated factors among school-age children in Dara Mallo and Uba Debretsehay districts, Southern Ethiopia. *Malar J*. 2020;19(1):369.
30. Inungu JN, Ankiba N, Minelli M, Mumford V, Bolekela D, Mukoso B, et al. Use of insecticide-treated mosquito net among pregnant women and guardians of children under five in the Democratic Republic of the Congo. *Malaria Research and Treatment*. 2017;2017.
31. Min KT, Maung TM, Oo MM, Oo T, Lin Z, Thi A, et al. Utilization of insecticide-treated bed nets and care-seeking for fever and its associated socio-demographic and geographical factors among under-five children in different regions: evidence from the Myanmar Demographic and Health Survey, 2015–2016. *Malaria Journal*. 2020;19(1):1-12.
32. Kanyangarara M, Hamapumbu H, Mamini E, Lupiya J, Stevenson JC, Mharakurwa S, et al. Malaria knowledge and bed net use in three transmission settings in southern Africa. *Malaria journal*. 2018;17(1):1-12.
33. Matovu SN, Wallhagen MI. Perceived caregiver stress, coping, and quality of life of older Ugandan grandparent-caregivers. *Journal of Cross-Cultural Gerontology*. 2020;35(3):311-28.
34. Andinda M, Mulogo E, Turyakira E, Batwala V. Predictors of sleeping under cost-free mosquito bed nets among children under-five years in Mbarara, Uganda: a household survey. *African health sciences*. 2019;19(1):1353-60.
35. Israel OK, Fawole OI, Adebawale AS, Ajayi IO, Yusuf OB, Oladimeji A, et al. Caregivers' knowledge and utilization of long-lasting insecticidal nets among under-five children in Osun State, Southwest, Nigeria. *Malaria journal*. 2018;17(1):1-9.

36. Scates SS, Finn TP, Wisniewski J, Dadi D, Mandike R, Khamis M, et al. Costs of insecticide-treated bed net distribution systems in sub-Saharan Africa. *Malaria Journal*. 2020;19(1):1-18.
37. Mohamed MH, Mohamed H, Hassan A. Factors Affecting Utilisation of Insecticide Treated Net among Household with Children Less than Five Years in Hodan District, Mogadishu, Somalia. *Journal of Biosciences and Medicines*. 2020;8(08):194.
38. Alemu MB, Asnake MA, Lemma MY, Melak MF, Yenit MK. Utilization of insecticide treated bed net and associated factors among households of Kola Diba town, North Gondar, Amhara region, Ethiopia. *BMC research notes*. 2018;11(1):1-5.
39. Babalola S, Adedokun ST, McCartney-Melstad A, Okoh M, Asa S, Tweedie I, et al. Factors associated with caregivers' consistency of use of bed nets in Nigeria: a multilevel multinomial analysis of survey data. *Malar J*. 2018;17(1):280.
40. Yirsaw AN, Gebremariam RB, Getnet WA, Mihret MS. Insecticide-treated net utilization and associated factors among pregnant women and under-five children in East Belessa District, Northwest Ethiopia: using the Health Belief model. *Malaria Journal*. 2021;20(1):1-12.
41. Koroma JM, Wang Y, Guo X, Zhang X, Ngobeh JJK, Gabir AMEA, et al. A Cross-Sectional Survey on Knowledge, Attitudes and Practices Towards Malaria Control and Prevention Among Caregivers of Children Under-5 in the Western Area of Sierra Leone. 2021.
42. Mubarick Nunbaso A, Assibi Ziba F, Nimota Nukpezah R, Abdulai A, Ayisha Evelyn M, Zakiyya M, et al. Awareness and Use of Insecticide-Treated Mosquito Bed Nets (ITNs) among Students in the Second Cycle Institutions in the Tamale Metropolis of Northern Region of Ghana. 2021.
43. Jumbam DT, Stevenson JC, Matoba J, Grieco JP, Ahern LN, Hamainza B, et al. Knowledge, attitudes and practices assessment of malaria interventions in rural Zambia. *BMC public health*. 2020;20(1):1-15.
44. Woday A, Mohammed A, Gebre A, Urmale K. Prevalence and associated factors of malaria among febrile children in Afar region, Ethiopia: a health facility based study. *Ethiopian journal of health sciences*. 2019;29(5).
45. Ricotta E, Oppong S, Yukich JO, Briët OJ. Determinants of bed net use conditional on access in population surveys in Ghana. *Malaria journal*. 2019;18(1):1-9.
46. Mekuria M, Binegde DN, Derega J, Teferi Bala E, Tesfa B, Deriba BS. Insecticide-Treated Bed Net Utilization and Associated Factors Among Households in Ilu Galan District, Oromia Region, Ethiopia. *Environmental Health Insights*. 2022;16:11786302221078122.
47. Carnevale P, Gay F. Insecticide-treated mosquito nets. *Malaria Control and Elimination: Springer*; 2019. p. 221-32.
48. Finda MF, Moshi IR, Monroe A, Limwagu AJ, Nyoni AP, Swai JK, et al. Linking human behaviours and malaria vector biting risk in south-eastern Tanzania. *PloS one*. 2019;14(6):e0217414.
49. Boadu I, Nsemani W, Ubachukwu P, Okafor F. Knowledge and prevalence of malaria among rural households in Ghana. *Journal of Community Medicine Health Education*. 2020;10(673):2.
50. Konlan K, Japiong M, DodamKonlan K, Afaya A, Salia S, Kombat J. Utilization of Insecticide Treated Bed Nets (ITNs) among Caregivers of Children under Five Years in the Ho Municipality. *Interdisciplinary Perspectives on Infectious Diseases*. 2019;2019.
51. Perkins JM, Krezanoski P, Takada S, Kakuhihire B, Batwala V, Tsai AC, et al. Social norms, misperceptions, and mosquito net use: a population-based, cross-sectional study in rural Uganda. *Malaria journal*. 2019;18(1):1-13.

## 12. Annexes

Annex 1. English version of participant's consent form

Questionnaires Insecticide Treated Bed Nets (ITNs) utilization and associated factors among under- five years old children in Lasta district, North Wollo zone Amhara, Ethiopia, 2022.

Questionnaire ID Number\_\_\_\_\_

District Name \_\_\_\_\_ Kebele name\_\_\_\_\_

### Section.1.GeneralInformation

Hello. My name is ----- and I am part of a team of people who are carrying out a survey on “Assessment of utilization of ITNs and association factor among households with under five children for malaria control in Lasta District” (Show a letter of approval from DHO). We would very much appreciate your participation in this survey. We would like to ask you some questions and it will take about 25 minutes. Your answers will remain confidential, and we will not be taking down your name or address, so your answers will be anonymous. Participation in this survey is voluntary and you can choose not to answer any individual question or all of the questions. However, we hope that you will participate in this survey since your views are important. At this time, do you want to ask me anything about the survey? May I begin the interview now? Start time\_\_\_\_ End time\_\_\_\_ Date\_\_\_\_/\_\_\_\_/\_\_\_\_ Respondent agrees to be interviewed.....Care taker does not agree to be interviewed-----  
END

Annex 2. English version of participant's questionnaires

| No_                                 | Question                         | Response  | Skip |
|-------------------------------------|----------------------------------|---|------|
| <b>I. Socio-demographic factors</b> |                                  |   |      |
| 101                                 | Age of care taker                | -----in years   |      |
| 102                                 | sex of care taker                | 1) Male<br>2) Female  |      |
| 103                                 | Educational status of care taker | 1)Unable to read and write<br>2)Able to read and write<br>3)Primary<br>4 )Secondary<br>5)above                  |      |
| 104                                 | Occupation                       | 1)Farmer<br>2)House wife<br>3)Student<br>4)Unemployed<br>6)Daily laborer<br>7)Government employee<br>8)Merchant |      |
| 105                                 | What is your marital status?     | 1) Married<br>2) Single<br>3) Divorced<br>4)Widowed   |      |
| 106                                 | Religion                         | 1) Orthodox<br>2)Muslim   |      |

|     |  |  |         |
|-----|--|--|---------|
| II  | <b>Intra-household factors</b>   |  |         |
| 201 | How many children below 5 years live this household?                   | 1) One<br>2) Two<br>3) Three<br>4) More than three                     |         |
| 202 | Number of people live in this household?                               | -----  |         |
| 203 | How many sleeping spaces are there in this household?                  | 1) one<br>2)Two<br>3) Three<br>4) Four<br>5) Five<br>6) More than five | Observe |
| 204 | How many sleeping rooms were ITNS hanged?                              | _____  | observe |
| III | <b>Knowledge related factors</b>                                       |  |         |
| 301 | Do you know how to hang the ITN  | 1)Yes<br>2)No  |         |
| 302 | Do you know malaria transmission                                       | 1)Yes<br>2)No  |         |
| 303 | If yes, what are the main Transmition                                  | -----  |         |
| 304 | do you know sign an symptom of malaria                                 | 1)Yes<br>2)No  |         |
| 305 | Have you Ever heard about ITNs use                                     | 1)Yes<br>2)No  |         |
| 306 | Do you know ITNs are necessary in preventing malaria among under-fives | 1)Yes<br>2)No  |         |



|     |  |   |                   |
|-----|--|---|-------------------|
|     | year   |   |                   |
| 307 | Do you think that malaria is caused other than mosquitoes bite                                   | 1)Yes<br>2)No   |                   |
| 308 | Do you know the use of treated and Non treated ITN   | 1)Yes<br>2)No   |                   |
| 309 | Do you think that social support network determines bed nets use for children under- five years? | 1) Yes<br>2) No   |                   |
| 310 | Have you ever been you and your partner discusses about ITN use for children under five years?   | 1)Yes<br>2)No   |                   |
| IV  | <b>ITN related factors</b>   |   |                   |
| 401 | Do you have ITNs in the house  | 1.yes<br>2.No   | If No, skip Q.402 |
| 402 | If yes, How many ITNs do you have?   | 1)One<br>2)Two<br>3)Three<br>4)above three                          |                   |
| 403 | What type of ITN do you have?  | 1)Treated ITN<br>2)Non-treated ITN                                  |                   |
| 404 | Source of ITN  | 1) Support from any organization<br>2)Support from Family<br>3)Paid |                   |
| 405 | Utilization of ITN in under five   | 1)Yes<br>2)No   |                   |

Annex 3.Checklist for direct observation

| S.No | Category                          | Response           |
|------|-----------------------------------|--------------------|
| 405  | Number of beds or places of sleep | 1)One<br><br>2)Two |

|     |  |  |
|-----|--|--|
|     |  | 3)Three and above                        |
| 406 | Number of beds /places of sleep observed with bed nets               | 1)One<br>2)Two<br>3)Three and above      |
| 407 | Number of bed nets observed in the household                         | 1) One<br>2) Two<br>3)Three and above    |
| 408 | The type of bed net that household owned                             | 1)Re- treatable<br>2)Permanently treated |
| 409 | Is the bed net hanged(placed) properly over the bed or sleeping area | 1)Yes<br>2)No                            |
| 410 | Is there any hole(throne) in the bed net                             | 1)yes<br>2)No                            |

Annex 4.Amharic version of participant's consent form

አባሪ 1.የተሳታፊ ስምምነት ቅጽ

የወባ በሽታ መከላከያ በሆነው በአልጋ አጎበሮ አጠቃቀም እና ተዛማጅ ምክንያቶች ዙሪያ ከአምስት ዓመት በታች ለሆኑ ሕፃናት ቤተሰቦች የተዘጋጀ መጠይቅ በላስታ ወረዳ፡ ሰሜን ወሎ ዞን፤ አማራ ፣ ኢትዮጵያ ፤ 2021

የተሳታፊ መታወቂያ ቁጥር \_\_\_\_\_ የወረዳ ስም \_\_\_\_\_ የቀበሌ ስም \_\_\_\_\_

ክፍል. 1. አጠቃላይ መረጃ

ሰላም። ስሜ -----ይባላል። እና እኔ የወባ በሽታን ለመቆጣጠር ከአምስት ዓመት በታች በሆኑ ህፃናት ባላቸው ቤተሰቦች መካከል የአልጋ አጎበሮ አጠቃቀም ላይ ጥናት ለማድረግ የመጣሁ ነኝ። በላስታ ወረዳ” (ከ ጤና ጽ/ቤት የድጋፍ ደብዳቤን ያሳዩ)። በዚህ የዳሰሳ ጥናት ውስጥ ተሳትፎዎን በጣም እናደንቃለን። አንዳንድ ጥያቄዎችን ልጠይቅዎት እና ወደ 25 ደቂቃዎች ያህል ይወስዳል። መልሶችዎ ሚስጥራዊ ሆነው ይቆያሉ ፣ እና እኔ ስምዎን ወይም አድራሻዎን አንፈልግም ፣ ስለዚህ መልሶችዎ ስም -አልባ ይሆናሉ።

በዚህ የዳሰሳ ጥናት ውስጥ መሳተፍ በፈቃደኝነት ላይ የተመሠረተ ሲሆን ማንኛውንም የግለሰብ ጥያቄ ወይም ሁሉንም ጥያቄዎች ላለመመለስ መምረጥ ይችላሉ። ሆኖም ፣ የእርስዎ አስተያየት አስፈላጊ ስለሆነ በዚህ የዳሰሳ ጥናት ውስጥ እንደሚሳተፉ ተስፋ እናደርጋለን።

በዚህ ጊዜ ስለ ዳሰሳ ጥናቱ ማንኛውንም ነገር ሊጠይቁኝ ይፈልጋሉ? አሁን ቃለ መጠይቁን ልጀምር? የመነሻ ሰዓት \_\_\_\_\_ የማብቂያ ጊዜ \_\_\_\_\_ ቀን \_\_\_\_/\_\_\_\_/\_\_\_\_ ተጠሪ ቃለ መጠይቅ ለማድረግ ተስማማ .....

ተጠሪ በቃለ መጠይቅ አይስማማም ----- መጨረሻ።

| ተፎታ         | መጠይቅ                    | ምላሽ   | ዝላል |
|-------------|-------------------------|---|-----|
| ስነ ህዝባዊ መረጃ |                         |   |     |
| 101         | የአሳድጊ ወይም የወላጅ እድሜ      | -----በአመት   |     |
| 102         | የአሳድጊ ወይም የወላጅ ፆታ       | 1) ወንድ<br>2) ሴት   |     |
| 103         | የአሳድጊ ወይም የወላጅ የት/ት ደረጃ | 1)መፃፍና ማንበብ የማይችል<br>2)መፃፍና መንበብ የሚችል<br>3)አንድኛ ደረጃ<br>4 )ሁለተኛ ደረጃ<br>5)ከሁለተኛ ደረጃ በላይ |     |
| 104         | የስራ ድርሻ                 | 1)አርሶ አደር<br>2)የቤት እመቤት<br>3)ተማሪ<br>4)ስራ አጥ<br>6)ቀን ሰራተኛ<br>7)የመንግስት ሰራተኛ<br>8)ነጋዴ    |     |

|     |                            |   |  |
|-----|----------------------------|---|--|
| 105 | የጋብቻ ሁኔታ ?                 | 1) ያገባ<br>2) ያላገባ<br>3) የፈታ<br>4) የሞተባት/በት                      |  |
| 106 | ሃይማኖት                      | 1) ኦርቶዶክስ ተዋህዶ<br>2) ሙስሊም                                       |  |
| II  | <b>የቤት ውስጥ ባህርያት መጠይቆች</b> |   |  |
| 201 | ስንት ከአምስት አመት በታች ህፃናት አሉ? | 1) አንድ<br>2) ሁለት<br>3) ሶስት<br>4) ከሶስት በላይ                       |  |
| 202 | በቤቱ ውስጥ ስንት ሰው ይኖራል ?      | -----   |  |
| 203 | ስንት የመኝታ ቦታዎች ወይም ክፍሎች አሉ? | 1) አንድ<br>2) ሁለት<br>3) ሶስት<br>4) አራት<br>5) አምስት<br>6) ከአምስት በላይ |  |

|     |  |               |  |
|-----|--|---------------|--|
| 204 | ከስንቱ የመኝታ ክፍሎች አገበር በአግባቡ ተሰቅሏል?         | _____         |  |
| III | ስነ ባህሪያዊ መረጃዎች                           |               |  |
|     | <b>የአሳዳጊ ወይም ተንከባካቢ እውቀት መረጃ</b>         |               |  |
| 301 | የአልጋ አገበር እንደት እንደሚሰቀል አውቃሉ?             | 1)አዎ<br>2)የለም |  |
| 302 | የወባ መከላከያ መንገዶችን ያውቃሉ?                   | 1)አዎ<br>2)የለም |  |
| 303 | ካወቁ ምን ምን ናቸው                            | -----         |  |
| 304 | የወባ በሽታ ምልክቶችን ያውቃሉ?                     | 1)አዎ<br>2)የለም |  |
| 305 | ስለ አልጋ አገበር ሰምተው ያውቃሉ?                   | 1)አዎ<br>2)የለም |  |
| 306 | የአልጋ አገበር የወባ ትንኝን ይከላከላል ብለው ያስባሉ?      | 1)አዎ<br>2)የለም |  |
| 307 | የወባ በሽታ በወባ ትንኝ አማካኝነት የሚመጣ ነው ብለው ያስባሉ? | 1)አዎ<br>2)የለም |  |
| 308 | የታከመ የአልጋ አገበርና ያልታከመ አገበር ጥቅምን ያውቃሉ?    | 1)አዎ<br>2)የለም |  |
|     | <b>የአሳዳጊዎች ወይም ተንከባካቢዎች አመለካከት</b>       |               |  |

|     |  |  |  |
|-----|--|--|--|
| 313 | ህፃናትን አጎበርን በእግባቡ እንድጠቀሙ የእርሳዎን እርዳታ ይፈልጋሉ ብለው ያስባሉ? | 1)አዎ<br>2)የለም  |  |
| 314 | እርሳዎና ቤተሰብዎ ስለ አጎበርና አጠቃቀሙ ተመካክረው ያውቃሉ?              | 1)አዎ<br>2)የለም  |  |
| IV  | <b>አጎበር አጠቃቀምን በተመለከተ መጠይቆች</b>                      | 1)አዎ<br>2)የለም  |  |
| 401 | ከቤትዎ የአልጋ አጎበር አለዎት                                  | 1)አዎ<br>2)የለም  |  |
| 402 | ካልዎት ምን ያህል ነው?                                      | 1) አንድ<br><br>2) ሁለት<br><br>3)ሶስት<br><br>4) ከሶስት በላይ |  |
| 403 | ምን አይነት የአልጋ አጎበር አለዎት ?                             | 1.የታከመ<br>2.ያልታከመ                                    |  |
| 404 | ከየት አገኙት   | 1. ከመንግስት<br>2.ከቤተሰብ<br>3.በግዥ                        |  |
| 405 | ከአምስት አመት በታች ያለ ህፃን አጎበር ተጠቅሟል?                     | 1.አዎ<br>2.የለም  |  |

Annex 3. የምልከታ መጠይቆች

| S.No | ጥያቄ                                    | ምላሽ                                  |
|------|--|--------------------------------------|
| 405  | የመኝታ ክፍሎች ወይም ቦታዎች                     | 1)አንድ<br>2)ሁለት<br>3)ሶስትና ከዚያ በላይ     |
| 406  | የአልጋ አጎበር የተሰቀለባቸው የመኝታ ቦታዎች ወይም መኝታዎች | 1)አንድ<br>2)ሁለት<br>3)ሶስትና ከዚያ በላይ     |
| 407  | በቤት ውስጥ ያለ ጠቅላላ አጎበር                   | 1) አንድ<br>2) ሁለት<br>3)ሶስትና ከዚያ በላይ \ |
| 408  | የአልጋ አጎበሩ ዐይነት                         | 1)ሊታከም የሚችል<br>2)የማይታከም              |
| 409  | የአልጋው አጎበር በትክክል ተሰቅሏል ወይ              | 1)አዎ<br>2)የለም                        |
| 410  | የአልጋ አጎበሩ የተቀደደ ነገር አለው ወይ             | 1)አዎ<br>2)የለም                        |





ባሕር ዳር ዩኒቨርሲቲ  
 ሕክምና እና ጤና ሳይንስ ኮሌጅ  
 የሥነ - ምግባር ገምጋሚ ቦርድ  
 ባሕር ዳር ፣ ኢትዮጵያ

Bahir Dar University  
 College of Medicine and Health Sciences  
 Institutional Review Board  
 Bahir Dar, Ethiopia



☒ ----- ☎ 058 89 99 275 /058 89 99271  
 Fax: ----- Website: www.bdu.edu.et/cmhs

E-mail :cmhs@bdu.edu.et  
 Facebook: College of Medicine and Health  
 Sciences, Bahir Dar University, Ethiopia

### IRB's Decision

Meeting No.: fullreview /2022  
 Protocol number: 363/2021

Date: March 11, 2022  
 assigned No: 003

|   |  |
|---|--|
| Protocol Title: - Assessment of insecticide treated bed nets utilization and associated factors among under five years children in Lasta district, North wollo zone , Amhara region, north east ethiopia,2021 |  |
| Principal investigator:   | Asfaw Hailu zerihun  |
| Co-investigators  |  |
| Institute:  | College of Medicine and Health Sciences, Bahir Dar University                                      |
| Elements Reviewed (CMHS/IRB 01 - 008)   | <input checked="" type="checkbox"/> Attached <input type="checkbox"/> Not attached                 |
| Review of Revised Application<br><input type="checkbox"/> Yes <input checked="" type="checkbox"/> No  | Date of Previous review:   |
| Decision of the meeting:  | <input checked="" type="checkbox"/> Approved <input type="checkbox"/> Approved with Recommendation |
|   | <input type="checkbox"/> Resubmission <input type="checkbox"/> Disapproved                         |

Elements approved:

1. Protocol Version No.: 01
2. Protocol Version Date March 11, 2022
3. Informed Consent Version: 01
4. Informed Consent Version Date: March 11, 2022

Obligations of the PI:

- ✓ Comply with Standard National and International Ethical Guidelines
- ✓ All Amendments and Changes made in the Protocol and Consent Needs IRB Approval
- ✓ Report SAE within 10 days of the event
- ✓ End of the study, including manuscript and thesis works should be reported to IRB

To NRERC ☒

Institutional Review Board (IRB) Approval: Period from 03/11/2022 to 06/11/2022

Follow-up Report Expected in:

3months ☒ 6months ☐ 9months ☐ One Year ☐

Chairperson, IRB : Mulusew Andualem(PhD)

Signature

Mulusew Andualem Asemahagn  
 (PhD)  
 Director of Research and  
 Management Office

Date: March 11, 2022



## 1. DECLARATION AND APPROVAL SHEET

### Declaration

This is to certify that the thesis entitled "insecticide treated bed nets utilization and associated factors among under five children in Lasta districts, North Wollo zone, Amhara, Ethiopia" submitted in partial fulfillment of the requirements for the degree of Masters of Public Health in Field Epidemiology of Departments of Epidemiology and Biostatistics, Bahir Dar University, is a record of original work carried out by me and has never been submitted to this or any institution to get any other degree or certificates. The assistance and help I received during the course of this investigation have been duly acknowledged.

Asfaw Hailu  
Name of the student

Asfaw  
Signature

August 17, 2022  
Date

### Approval sheet of thesis for defense

I hereby certify that I have supervised, read, and evaluated this thesis titled "insecticide treated bed nets utilization and associated factors among under five children in Lasta districts, North Wollo zone, Amhara, Ethiopia" by Asfaw Hailu prepared under my guidance. I recommend the thesis be submitted for oral defense

Kekadnew Muluat  
Advisor's name

Signature

Date

12-12-2014 E.C

Gebreyes W  
Co-Advisor's name

Signature

Date

12-12-2014 E.C

Behiyaw Woldu  
Department Head

Signature

Date

12-12-2014 E.C



**Approval of thesis for defense result**

As members of the board of examiners, we examined this thesis entitled "insecticide treated bed nets utilization and associated factors among under five children in Lasta districts, North Wollo zone, Amhara, Ethiopia" by Asfaw Hailu. We hereby certify that the thesis is accepted for fulfilling the requirements for the award of the degree of "Masters of Public Health in Field Epidemiology".

**Board of Examiners**

| External examiner name | Signature | Date            |
|------------------------|-----------|-----------------|
| _____                  | _____     | _____           |
| _____                  | _____     | _____           |
| Internal examiner name | Signature | Date            |
| <u>Abbeaw Gedef</u>    | <u>Ng</u> | <u>12/12/14</u> |
| _____                  | _____     | _____           |
| Chair person's name    | Signature | Date            |
| _____                  | _____     | _____           |